





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/11, 9/12, 15/85, C12Q 1/68		A2	(11) International Publication Number: WO 99/38964
			(43) International Publication Date: 5 August 1999 (05.08.99)
(21) International Application Number: PCT/GB99/00308 (22) International Filing Date: 29 January 1999 (29.01.99) (30) Priority Data: 9801902.9 29 January 1998 (29.01.98) GB (71) Applicant (for all designated States except US): CANCER RESEARCH CAMPAIGN TECHNOLOGY LIMITED [GB/GB]; Cambridge House, 6-10 Cambridge Terrace, Regent's Park, London NW1 4JL (GB). (72) Inventor; and (75) Inventor/Applicant (for US only): KEITH, William, Nicol [GB/GB]; CRC Dept. of Medical Oncology, University of Glasgow, CRC Beatson Laboratories, Garscube Estate, Switchback Road, Glasgow G61 1BD (GB). (74) Agents: CRIPPS, Joanna, E. et al.; Mewburn Ellis, York House, 23 Kingsway, London WC2B 6HP (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published Without international search report and to be republished upon receipt of that report.	

(54) Title: PROMOTER REGIONS OF THE MOUSE AND HUMAN TELOMERASE RNA COMPONENT GENES

(57) Abstract

The present invention relates to the identification of the genomic promoter region of the human and mouse telomerase RNA gene. Telomerase activity is necessary for the unrestricted proliferative capacity of many human cancers. It is proposed that mutation or dysregulation of the telomerase repression pathway may cause reactivation or upregulation of telomerase expression in cancer. The invention provides details of elements important for the regulation of telomerase RNA genes, including the Sp family of transcription factors. There is further provided methods for screening for elements having the ability for suppressing telomerase RNA gene promoter activity and use of such elements in the treatment of cancers. In addition, evidence is also provided for the development of new transcription based therapies for cancer and for genetic approaches to targeting therapeutic genes to cancer cells. Namely, (1) transcriptional repression and the disruption of signal transduction pathways regulating telomerase activation. (2) Tumour specific gene expression for genetic therapy via telomerase RNA gene promoters.

**A****B**

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

A gene promoterField of the invention

5 The present invention relates to the cloning of a genomic promoter region of the human and mouse telomerase RNA gene. Particularly, but not exclusively, it relates to the identity of elements important for the regulation of telomerase RNA genes.

10 Background of the invention

Telomeres are found at the end of linear chromosomes and consist of short repetitive sequences essential for the maintenance of normal chromosome structure and function (Wellinger & Sen, 1997). With each cell division, 15 telomeres shorten due to the inability of DNA polymerases to replicate the ends of linear DNA molecules. However, telomere erosion is counteracted by the activity of the enzyme telomerase, a ribonuclear protein with reverse transcriptase activity, which adds telomeric repeats to the chromosomal termini (Morin, 1997; Nakamura et al., 1997). 20 The genes for the human, (hTR), and mouse, (terc), RNA components have recently been cloned, as has the human protein component, (hTRT) (Blasco et al., 1995; Feng et al., 1995; Nakamura et al., 1997; Soder et al., 1997b; 25 Soder et al., 1997c). Whilst telomerase expression is detectable in normal embryonic tissues and germline stem cells, telomerase expression is repressed in most normal postnatal somatic cells (Blasco et al., 1995; Feng et al., 1995; Prowse & Greider, 1995; Soder et al., 1997a; Wright et al., 1996). The lack of telomerase expression may be the 30 major reason for the progressive loss of telomeric

sequences in somatic cells, which is considered to be one regulatory mechanism which monitors the number of times a cell divides before entering replicative senescence (Campisi, 1997). However, although telomerase appears to be stringently repressed in normal somatic tissues, there is substantial evidence to suggest that telomerase is expressed in the majority of human cancers and contributing to the immortal phenotype through the maintenance of telomere integrity (Holt et al., 1997; Kim, 1997; Shay & Bacchetti, 1997).

The regulation of telomerase expression is a complex issue including transcriptional activity of the telomerase RNA and protein component genes, interaction of telomerase with other telomere associated proteins and post-translational modification of the enzyme complex. However, at present there are few studies which directly address the mechanisms regulating telomerase expression in normal and cancer cells (Bodnar et al., 1996; Broccoli et al., 1997; Li et al., 1997; Mandal & Kumar, 1997; Morin, 1997; Nakamura et al., 1997; Soder et al., 1997a).

Studies aimed at relating genome stability to human cellular senescence have recently placed considerable emphasis on telomerase expression as a central unifying mechanism underlying the immortal phenotype of many cancers (Breslow et al., 1997). The absence of telomerase activity from normal somatic cells has led to the proposal that telomere shortening may be a molecular clock which contributes to the onset of cellular senescence in normal cells (Harley & Villeponteau, 1995; Holt et al., 1996). Conversely, the reactivation or expression of telomerase may be a major mechanism by which cancer overcome normal

cellular senescence (Kim et al., 1994; Parkinson et al., 1997). Information on telomerase activity in tumours almost exclusively derives from the *in vitro* telomere repeat amplification protocol, (TRAP), and these have shown that
5 telomerase activity may be present in greater than 80% of tumour biopsies yet absent or reduced in normal somatic tissue (Breslow et al., 1997; Kim et al., 1994; Raymond et al., 1996; Shay & Wright, 1996). However, TRAP assay alone will not reveal the true complexities of telomerase
10 regulation, and it is generally recognised that a number of molecular approaches will be required to understand telomere length regulation and telomerase activity (Breslow et al., 1997; Holt et al., 1996; Lundblad & Wright, 1996; Parkinson et al., 1997; Raymond et al., 1996; Soder et al.,
15 1997b). Recently, the present inventors and others have introduced a number of more direct *in situ* approaches to study the telomerase RNA gene, (hTR) and its expression in tumours (Soder et al., 1997; Yashima et al., 1997b).

US patent 5583016 (Geron Corp.) Discloses a 2.4kb
20 sequence of the Telomerase RNA gene. However, there is no disclosure of the promoter elements or provision of functional evidence to show the promoter is active.

Summary of the invention

25 The levels of telomerase RNA gene expression vary during normal development and between normal and cancerous cells and tissues (Avilion et al., 1996; Bestilny et al., 1996; Blasco et al., 1995; Blasco et al., 1996; Bodnar et al., 1996; Broccoli et al., 1996; Feng et al., 1995;
30 Kuniyasu et al., 1997; Soder et al., 1997a). The present inventors have appreciated that knowledge of telomerase RNA

gene expression should aid understanding of the signal transduction pathways linking telomere attrition to proliferation, cellular senescence, differentiation and oncogenesis. As a first step towards this goal, they have
5 cloned the promoter regions of the human, (hTR), and mouse, (terc), telomerase RNA genes in order to identify the regulatory elements controlling telomerase RNA gene transcription. Further, the present inventors have investigated the possible levels of telomerase regulation
10 in vivo.

These studies have a number of implications for the development of new transcription based therapies for cancer (Cai et al., 1996; Connors, 1995; Miller & Whelan, 1997; Peterson & Baichwal, 1993). Directly down-regulating
15 expression of the telomerase RNA gene through manipulation of transcription factors should be an effective anticancer therapy and the cloning of the hTR gene promoter will allow the analysis of therapeutic molecules which modulate hTR promoter activity (Cai et al., 1996; Peterson & Baichwal,
20 1993; Sharma et al., 1997). Indeed, by using a human cell line which has telomerase activity, (HeLa) and one which expresses the hTR gene but is telomerase negative, (GM847), (Bryan et al., 1997), the present inventors provide a system in which the specificity of transcriptional
25 manipulation of hTR may be examined. In comparison to HeLa, the growth of GM847 does not appear to be dependant on telomerase expression, thus transcriptional targeting of hTR in GM847 should have no cellular effects, whereas HeLa should be sensitive to the predicted anti-proliferative
30 effects of the transcriptional targeting. In addition, the present inventors show tumour-specific patterns of hTR gene

expression with clear differentials in expression between cancerous and adjacent normal tissue, (Soder et al., 1997a (not yet published); Soder et al., 1997b).

5 Broadly, the present invention provides materials and methods relating to the telomerase RNA (TR) gene promoter and its effects in tumour development.

According to a first aspect there is provided a nucleic acid molecule comprising the TR gene promoter.

10 In a second aspect there is provided a nucleic acid molecule comprising a human TR gene promoter, preferably the promoter comprising a sequence of nucleotides shown in Fig. 4a. The promoter may comprise one or more fragments of the sequence shown in Fig. 4a, sufficient to promote gene expression. In particular, it may comprise or consist
15 essentially of a sequence of nucleotides extending at least 200 base pairs (bp), or 250 bp, 272 bp, or 300 bp, preferably at least 320 bp, more preferably at least 340 bp, even more preferably at least 400 bp upstream of the transcription start site. More preferably, the promoter may
20 comprise a sequence of nucleotides of at least 230 bp in length starting at position -42 bp upstream of the transcription start site (Fig. 4a and Fig. 5a). Even more preferably the promoter may comprise a sequence of nucleotides of at least 230bp in length between positions
25 -272 bp and -42 bp (Fig. 4a and Fig. 5a).

In a further preferred form the promoter comprises or consists essentially of the construct designated hProm505 being a sequence of nucleotides of approximately 505 bp in length from position -463 as shown in Fig. 4 or the
30 construct designated hProm867 being 867 bp in length from position -798 bp (Fig. 4a and Fig. 5a).

In a third aspect of the present invention there is provided a nucleic acid molecule comprising mouse Telomerase RNA (*terc*) gene promoter, preferably comprising or consisting essentially of the sequence as shown in Fig. 4b. In a preferred form, the present invention provides a nucleic acid molecule comprising a *terc* promoter, said promoter comprising or consisting essentially of a nucleotide sequence extending at least 94 bp, preferably at least 100 bp, more preferably at least 120 bp, even more preferably at least 150 bp upstream of the transcription start site. Preferably the promoter comprises or consists essentially of a nucleotide sequence of approximately 73 bp in length between -94 bp and -22 bp as shown in Fig. 4b and Fig. 5b.

In further preferred form, the present invention provides a nucleic acid molecule comprising or consisting essentially of the construct designated mProm208 or the construct designated mProm628 as shown in Fig. 4 or Fig. 5b.

An even smaller portion of the nucleotide sequences mentioned above may be used as long as the promoter activity is retained. Such nucleotide sequences may be fragments being 200 nucleotides or fewer in length (e.g. 150, 100, 50, 40, 35, 30, 25, or 20). Restriction enzymes or nucleases may be used to digest the nucleic acid, followed by the appropriate assay (for example as illustrated herein using luciferase constructs) to determine the minimal sequence required. A preferred embodiment of the present invention provides a nucleic acid molecule with the minimal nucleotide sequence shown in Fig. 4a or Fig. 4b required for promoter activity. The minimal

promoter is situated between -272 bp and -42 bp as shown in Fig. 4a and between -94 bp and -22 bp as shown in Fig. 4b.

The promoter may comprise one or more sequence motifs or elements conferring developmental and/or tissue-specific regulatory control or expression. Other regulatory sequences may be included, for instance as identified by mutation or digest assay in an appropriate expression system or by sequence comparison with available information, e.g. using a computer to search on-line databases.

By "promoter" is meant a sequence of nucleotides from which transcription may be initiated of DNA operably linked downstream (i.e. in the 3' direction on the sense strand of double-stranded DNA). In the present case, the sequence of nucleotides is derived from either hTR promoter region or the mouse *terc* promoter. In other words, the nucleotide sequence may be identical to that of the hTR or the *terc* promoter in both size and sequence, or it may have been modified by insertion, addition, deletion or substitution of one or more nucleotides without fundamentally altering the essential activity of the promoter as compared to the hTR or *terc* promoter activity respectively. The essential activity can be determined by analysis of individual promoter elements as to their size, sequence and susceptibility to transcriptional control elements as described herein. Further, within the meaning of the term "promoter" is a sequence of nucleotides derived from either hTR promoter region or the mouse *terc* promoter which on its own lacks the essential activity but which can regain activity on addition of further nucleotide sequence, particularly nucleotide sequence comprising further

promoter elements known to those skilled in the art.

"Operably linked" means joined as part of the same nucleic acid molecule, suitably positioned and orientated for transcription to be initiated from the promoter. DNA
5 operably linked to a promoter is "under transcriptional initiation regulation" of the promoter.

The present invention extends to a promoter which is an allele, mutant, variant or derivative, by way of nucleotide addition, substitution or deletion of a promoter
10 sequence as provided herein. Systematic or random mutagenesis of nucleic acid to make an alteration to the nucleotide sequence may be performed using any technique known to those skilled in the art. One or more alterations to a promoter sequence according to the present invention
15 may increase or decrease promoter activity, or increase or decrease the magnitude of the effect of a substance able to modulate the promoter activity.

"Promoter activity" is used to refer to ability to initiate transcription. The level of promoter activity is
20 quantifiable for instance by assessment of the amount of mRNA produced by transcription from the promoter or by assessment of the amount of protein product produced by translation of mRNA produced by transcription from the promoter. The amount of a specific mRNA present in an
25 expression system may be determined for example using specific oligonucleotides which are able to hybridise with the mRNA and which are labelled or may be used in a specific amplification reaction such as the polymerase chain reaction. Use of a reporter gene as discussed
30 further below facilitates determination of promoter activity by reference to protein production.

In various embodiments of the present invention a promoter which has a sequence that is a fragment, mutant, allele, derivative or variant, by way of addition, insertion, deletion or substitution of one or more nucleotides, of the sequence of the hTR promoter shown in Fig. 4a or the *terc* promoter shown in Fig. 4b, has at least about 60% homology with one or both of the shown sequences, preferably at least about 70% homology, more preferably at least about 80% homology, more preferably at least about 90% homology, more preferably at least about 95% homology. Such homology may be found over a sequence of at least 10 nucleotides, preferably of at least 20 nucleotides, more preferably of at least 30 nucleotides and even more preferably of at least 40 nucleotides. Such fragments themselves individually represent aspects of the present invention.

The sequence in accordance with an embodiment of the invention may hybridise with one or both of the shown sequences, or the complementary sequences (since DNA is generally double-stranded). The sequence may have the ability to promote transcription (i.e. have "promoter activity") in normal embryonic tissues and germline stem cells.

In accordance with the present invention, nucleic acids having the appropriate level of sequence homology with the respective promoter regions may be identified by using hybridization and washing conditions of appropriate stringency. For example, hybridizations may be performed, according to the method of Sambrook et al., (22) using a hybridization solution comprising: 5X SSC, 5X Denhardt's reagent, 0.5-1.0% SDS, 100 µg/ml denatured, fragmented

salmon sperm DNA, 0.05% sodium pyrophosphate and up to 50% formamide. Hybridization is carried out at 37-42°C for at least six hours. Following hybridization, filters are washed as follows: (1) 5 minutes at room temperature in 2X SSC and 1% SDS; (2) 15 minutes at room temperature in 2X SSC and 0.1% SDS; (3) 30 minutes-1 hour at 37°C in 1X SSC and 1% SDS; (4) 2 hours at 42-65°C in 1X SSC and 1% SDS, changing the solution every 30 minutes.

One common formula for calculating the stringency conditions required to achieve hybridization between nucleic acid molecules of a specified sequence homology is (Sambrook et al., 1989):

$$T_m = 81.5^{\circ}\text{C} + 16.6\text{Log} [\text{Na}^+] + 0.41(\% \text{ G+C}) - 0.63 (\% \text{ formamide}) - 600/\text{\#bp in duplex}$$

As an illustration of the above formula, using $[\text{Na}^+] = [0.368]$ and 50% formamide, with GC content of 42% and an average probe size of 200 bases, the T_m is 57°C. The T_m of a DNA duplex decreases by 1 - 1.5°C with every 1% decrease in homology. Thus, targets with greater than about 75% sequence identity would be observed using a hybridization temperature of 42°C. Such a sequence would be considered substantially homologous to the respective promoter sequences of the present invention.

On the basis of the nucleotide sequences given herein (Fig. 4a and Fig. 4b), oligonucleotide probes or primers may be designed. Generally specific primers are upwards of 14 nucleotides in length but not more than 18 to 20. Those skilled in the art are well versed in the design of primers for use processes such as PCR. Primer sequences which themselves individually form part of the present invention are given in Fig. 6.

Further provided by the present invention is a nucleic acid construct comprising a TR promoter region or a fragment, mutant, allele, derivative or variant thereof able to promote transcription, operably linked to a heterologous gene, e.g. a coding sequence. By "heterologous" is meant a gene other than TR gene. Modified forms of TR are generally excluded. Generally, the gene may be transcribed into mRNA which may be translated into a peptide or polypeptide product which may be detected and preferably quantitated following expression. A gene whose encoded product may be assayed following expression is termed a "reporter gene", i.e. a gene which "reports" on promoter activity.

The reporter gene preferably encodes an enzyme which catalyses a reaction which produces a detectable signal, preferably a visually detectable signal, such as a coloured product. Many examples are known, including β -galactosidase and luciferase. β -galactosidase activity may be assayed by production of blue colour on substrate, the assay being by eye or by use of a spectrophotometer to measure absorbance. Luminescence, for example that produced as a result of luciferase activity, may be quantitated using a spectrophotometer. Radioactive assays may be used, for instance using chloramphenicol acetyltransferase, which may also be used in non-radioactive assays. The presence and/or amount of gene product resulting from expression from the reporter gene may be determined using a molecule able to bind the product, such as an antibody or fragment thereof. The binding molecule may be labelled directly or indirectly using any standard technique.

Those skilled in the art are well aware of a multitude of possible reporter genes and assay techniques which may be used to determine gene activity. Any suitable reporter/assay may be used and it should be appreciated that no particular choice is essential to or a limitation of the present invention.

Expression of a reporter gene from the promoter may be in an *in vitro* expression system or may be intracellular (*in vivo*). Expression generally requires the presence, in addition to the promoter which initiates transcription, a translational initiation region and transcriptional and translational termination regions. One or more introns may be present in the gene, along with mRNA processing signals (e.g. splice sites).

The present invention also provides a nucleic acid vector comprising a promoter as disclosed herein. Such a vector may comprise a suitably positioned restriction site or other means for insertion into the vector of a sequence heterologous to the promoter to be operably linked thereto.

Suitable vectors can be chosen or constructed containing appropriate regulatory sequences, including promoter sequences, terminator fragments, polyadenylation sequences, enhancer sequences, marker genes and other sequences as appropriate. For further details see, for example, *Molecular Cloning: a Laboratory Manual*: 2nd edition, Sambrook et al, 1989, Cold Spring Harbor Laboratory Press. Procedures for introducing DNA into cells depend on the host used, but are well known.

Thus, a further aspect of the present invention provides a host cell containing a nucleic acid construct comprising a promoter element, as disclosed herein,

operably linked to a heterologous nucleic acid coding sequence or a gene. In particular, the promotor of the present invention may be used to drive expression of an exogenous protein in selected cells. For example, the
5 coding sequence or gene may encode a cytotoxin which can then be selectively expressed in target cells e.g. tumour cells. Many such cytotoxins are known to the skilled person. They may include cytotoxins derived from viruses, bacteria or bacteriophages. Alternatively, the heterologous
10 gene may encode a endogenous protein and the promoter may be used to increase the expression of said endogenous protein in selective cells.

A still further aspect provides a method comprising introducing such a construct into a host cell. The
15 introduction may employ any available technique, including, for eukaryotic cells, calcium phosphate transfection, DEAE-Dextran transfection, electroporation, liposome-mediated transfection and transduction using retrovirus.

The introduction may be followed by causing or
20 allowing expression of the heterologous gene under the control of the promoter, e.g. by culturing host cells under conditions for expression of the gene.

In one embodiment, the construct comprising promoter and gene is integrated into the genome (e.g. chromosome) of
25 the host cell. Integration may be promoted by inclusion in the construct of sequences which promote recombination with the genome, in accordance with standard techniques.

Many known techniques and protocols for manipulation of nucleic acid, for example in preparation of nucleic acid
30 constructs, mutagenesis, sequencing, introduction of DNA into cells and gene expression, and analysis of proteins,

are described in detail in *Short Protocols in Molecular Biology*, Second Edition, Ausubel et al. eds., John Wiley & Sons, 1992, the disclosure of which is incorporated herein by reference.

5 Nucleic acid molecules, constructs and vectors according to the present invention may be provided isolated and/or purified (i.e. from their natural environment), in substantially pure or homogeneous form, free or substantially free of a TR coding sequence, or free or
10 substantially free of nucleic acid or genes of the species of interest or origin other than the promoter sequence. Nucleic acid according to the present invention may be wholly or partially synthetic. The term "isolate" encompasses all these possibilities.

15 Nucleic acid constructs comprising a promoter (as disclosed herein) and a heterologous gene (reporter) may be employed in screening for a substance able to modulate activity of the promoter. For therapeutic purposes, e.g. for treatment of cancers, a substance able to regulate the
20 promoter may be sought. A method of screening for ability of a substance to modulate activity of a TR promoter may comprise contacting an expression system, such as a host cell, containing a nucleic acid construct as herein disclosed with a test or candidate substance and
25 determining expression of the heterologous gene.

 The level of expression in the presence of the test substance may be compared with the level of expression in the absence of the test substance. A difference in expression in the presence of the test substance indicates
30 ability of the substance to modulate gene expression. A decrease in expression of the heterologous gene compared

with expression of another gene not linked to a promoter as disclosed herein indicates specificity of the substance for disrupting the TR promoter.

5 A promoter construct may be transfected into a cell line using any technique previously described to produce a stable cell line containing the reporter construct integrated into the genome. The cells may be grown and incubated with test compounds for varying times. The cells may be grown in 96 well plates to facilitate the analysis
10 of large numbers of compounds. The cells may then be washed and the reporter gene expression analysed. For some reporters, such as luciferase the cells will be lysed then analysed.

Constructs comprising one or more developmental and/or
15 time-specific regulatory motifs (as discussed) may be used to screen for a substance able to modulate the corresponding aspect of the promoter activity, e.g. cancer related expression.

There is interest in prolonging the cellular lifespan
20 of normal cells and keeping them normal rather than them aging or senescing. In this regard, the present invention provides the ability to upregulate or switch on at least TR and the protein component gene hTERT. This application may provide immortal normal cells for treating age related
25 diseases or to get unlimited amounts of commercially important normal cells. Alternatively, applications for substances able to suppress promoter activity are also provided, for example, for use in cancer therapy. Following identification of a substance which modulates or affects TR
30 gene promoter activity, the substance may be investigated further. Furthermore, it may be manufactured and/or used

in preparation, i.e. manufacture or formulation, of a composition such as a medicament, pharmaceutical composition or drug. These may be administered to individuals.

5 Thus, the present invention extends in various aspects not only to a substance identified using a nucleic acid molecule as a modulator of TR gene promoter activity, in accordance with what is disclosed herein, but also a pharmaceutical composition, medicament, drug or other
10 composition comprising such a substance, a method comprising administration of such a composition to a patient, e.g. for decreasing TR gene expression for instance in treatment of cancers, use of such a substance in manufacture of a composition for administration, e.g.
15 for decreasing TR gene expression for instance in treatment of cancers, and a method of making a pharmaceutical composition comprising admixing such a substance with a pharmaceutically acceptable excipient, vehicle or carrier, and optionally other ingredients.

20 Administration will preferably be in a "therapeutically effective amount", this being sufficient to show benefit to a patient. Such benefit may be at least amelioration of at least one symptom. The actual amount administered, and rate and time-course of administration,
25 will depend on the nature and severity of what is being treated. Prescription of treatment, eg decisions on dosage etc, is within the responsibility of general practitioners and other medical doctors.

30 A composition may be administered alone or in combination with other treatments, either simultaneously or sequentially dependent upon the condition to be treated.

Pharmaceutical compositions according to the present invention, and for use in accordance with the present invention, may comprise, in addition to active ingredient, a pharmaceutically acceptable excipient, carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material will depend on the route of administration, which may be oral, or by injection, e.g. cutaneous, subcutaneous or intravenous.

Pharmaceutical compositions for oral administration may be in tablet, capsule, powder or liquid form. A tablet may comprise a solid carrier such as gelatin or an adjuvant. Liquid pharmaceutical compositions generally comprise a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil or synthetic oil. Physiological saline solution, dextrose or other saccharide solution or glycols such as ethylene glycol, propylene glycol or polyethylene glycol may be included.

For intravenous, cutaneous or subcutaneous injection, or injection at the site of affliction, the active ingredient will be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability. Those of relevant skill in the art are well able to prepare suitable solutions using, for example, isotonic vehicles such as Sodium Chloride Injection, Ringer's Injection, Lactated Ringer's Injection. Preservatives, stabilisers, buffers, antioxidants and/or other additives may be included, as required.

Also included within the scope of the present

invention are substances that disrupt the ability of the TR gene promoter as herein described to regulate expression of the TR gene. These substances include any member that is capable of directly down-regulating telomerase RNA gene expression or which specifically blocks transcriptional activation of the TR gene promoters through interaction of the 5' regulatory sequences, for example, antisense oligonucleotides, transcription factors, synthetic oligonucleotides and peptide nucleic acids (hybrid molecules between peptides and nucleic acids) or factors that disrupt signal transduction pathways. Having identified the proximal region of the hTR promoter necessary to direct hTR transcription, a number of potential recognition sites for transcription factors have been identified by the present inventor (see Fig. 10). The present invention provides such transcription factors or functional fragments thereof for use in the modifying hTR transcription.

The present inventors have further devised reporter constructs, which form part of the invention, which can be introduced into cells or cell free systems to monitor promoter activity. This allows regulation of transcription and the development of small molecules specifically designed to disrupt the hTR transcriptional machinery. The reporter cell lines so produced also form part of the invention and are ideally suited to test any commercial telomerase inhibitors which may be evaluated, in combination with standard telomerase assays for specificity and mode of action. Therefore, the present invention provides methods and kits for testing telomerase inhibitors and molecules specifically designed to disrupt hTR

transcriptional machinery. Further, the present invention extends to the use of these reporter constructs to monitor promoter activity and regulate transcription.

5 The identification of the transcription factors regulating telomerase RNA gene expression and the elucidation of the specific molecules of the signal transduction pathways regulating expression will provide a wealth of targets for anticancer agents. Inhibitors may be developed by rational design based on the molecular
10 understanding of telomerase RNA gene expression or the screening of pharmaceutical compounds in screens based on promoter/reporter activity.

Therefore, the present invention provides a method of identifying transcription factors capable of regulating TR
15 gene expression comprising the steps of screening for compounds which disrupt TR gene promoter activity as determined by reporter activity. Telomerase inhibitors identified by such methods as described above are also within the scope of the present invention.

20 The work performed by the present inventors shows examples of clear differentials in hTR expression between cancerous and adjacent normal tissue which support the possibility of effective telomerase-based therapy (Soder et al 1997b; and Table 1 below). Indeed, the presence of high
25 levels of hTR expression in specific cancers shows that the hTR/terc promoter may be used for genetic therapies designed to target therapeutic genes to tumours, via tumour specific gene expression.

Therefore, the present invention provides the use of
30 nucleic acid molecules comprising promoter, promoter constructs or fragments thereof as described above for gene

therapy. This includes the preparation of a medicament for use in gene therapy.

5 Vectors such as viral vectors have been used in the prior art to introduce genes into a wide variety of different target cells. Typically the vectors are exposed to the target cells so that transfection can take place in a sufficient proportion of the cells to provide a useful therapeutic or prophylactic effect from the expression of the desired polypeptide. In the present case,
10 transcription and therefore expression of the desired polypeptide will be under the control of the promoter according to the present invention. The transfected nucleic acid comprising the promoter sequence and the sequence encoding the desired polypeptide may be permanently
15 incorporated into the genome of each of the targeted tumour cells, providing long lasting effect, or alternatively the treatment may have to be repeated periodically.

A variety of vectors, both viral vectors and plasmid vectors, are known in the art, see US Patent No. 5,252,479
20 and WO 93/07282. In particular, a number of viruses have been used as gene transfer vectors, including papovaviruses, such as SV40, vaccinia virus, herpesviruses, including HSV and EBV, and retroviruses. Many gene therapy protocols in the prior art have used disabled murine
25 retroviruses.

As an alternative to the use of viral vectors other known methods of introducing nucleic acid into cells includes electroporation, calcium phosphate co-precipitation, mechanical techniques such as
30 microinjection, transfer mediated by liposomes and direct DNA uptake and receptor-mediated DNA transfer.

Genetically directed enzyme prodrug therapy, (GDEPT), comprises two parts: a tumour specific promoter and the enzyme prodrug system. The properties of an optimal tumour selective transcriptional activation system can be summarized as, ideally tumour specific, only expressed in nonessential tissues, no cross-specificity with unusual but essential cell types, regulatory elements from gene cloned and sequenced, specific transcription factor binding sites identified, enhancer and inhibitory factors understood. The telomerase RNA gene promoters according to the present invention have these characteristics.

The present invention therefore provides the use of nucleic acid molecules comprising promoters as described herein to drive expression of enzyme-prodrug activation systems such as viral thymidine kinase and Gancyclovir, although many other systems known to those skilled in the art may also be used. Targeted gene expression via the telomerase RNA gene promoter may also be used in gene replacement strategies for cancer therapy. The present invention also provides a system for use in the control of neoplasia in a human or animal subject comprising a vector or other delivery system capable of selectively infecting tumour cells in said subject, said vector carrying a DNA or RNA sequence encoding an enzyme operably linked to a promoter sequence of the present invention, in association with a prodrug capable of being converted into an active compound by the action of said enzyme. The present invention further extends to a method of treating neoplasia in a human or animal subject requiring such treatment which comprises administering to the host an effective amount of a prodrug and a modified virus, said modified virus capable

of selectively infecting tumour cells in said subject, said virus carrying a DNA or RNA sequence encoding an enzyme capable of converting said prodrug to an active compound, said DNA or RNA sequence being operably linked to a promoter sequence in accordance with the present invention.

The present invention also extends to a method of treating neoplasia in a human or animal subject requiring such treatment which comprises administering to the host an effective amount of a prodrug and a non-viral vector system, said non-viral vector system capable of being selectively introduced into tumour cells in said subject, said vector system carrying a DNA or RNA sequence encoding an enzyme capable of converting said prodrug to an active compound, said DNA or RNA sequence being operably linked to a promoter in accordance with the present invention effective in expressing said enzyme in said cells. This aspect of the present invention is described in more detail below.

Genomic sequences encompassing the promoter region of hTR and *terc* have been cloned into green fluorescent protein (GFP) vectors and luciferase gene reporter vectors to assay for promoter activity, and deletion and mutation constructs have been developed. By introducing these constructs into primary cultures of keratinocytes/fibroblasts and neoplastic cell lines, which have varying levels of telomerase activity and telomerase RNA gene expression, genomic regions controlling repression or activation of hTR/*terc* expression may be identified. Human embryonic cells and murine embryonic stem cells may be used to study the developmental regulation of the expression.

The present invention also provides DNA binding assays for identifying transcription factors in crude cell extracts. As specific promoter regions have been identified by the present inventors, specific DNA-protein interactions may be analysed by gel mobility shift assay and DNase I footprinting which are well known techniques to those skilled in the art (Gene Transcription, eds Hames, B.D. and Higgins, S.J., IRL press, Oxford University Press, Oxford 1993). In addition, *in vitro* transcription assays for the hTR/terc promoters may be developed to study transcription factor interactions. Protein extracts from a) cell lines with and without telomerase RNA gene expression; b) early and late passage primary keratinocytes and fibroblasts; and c) tumour biopsies from patients, may be assayed for the involvement of previously characterised transcription factors by the use of purified transcription factors, oligonucleotide competition studies and antibodies to previously characterised factors. Should it be required, appropriate methods for the identification of the proteins responsible for the DNA-binding activity, such as UV crosslinking and protein purification and gene cloning may be performed. Proteins identified in this way will be considered as targets for inhibition and also form part of the invention.

The invention further provides a method for transcriptional analysis using transgenic mice. Transgenic mice with the human and mouse promoter/reporter constructs may be generated. Such transgenic mammals form part of the present invention. Also, the invention provides a method of creating transgenic mammals comprising the steps of injecting constructs into eggs of the mammal and re-

implanting the eggs into the female mammal. Ideally the mammal will be a mouse although those skilled in the art will readily contemplate the use of other mammals such as rats or rabbits. Mice carrying constructs in the germline may be used to establish transgenic lines. The invention also includes the use of human or mouse promoter/reporter constructs for the generation of transgenic mammals. The use of transgenic mice for the analysis of transcription factors has a number of advantages over tissue culture cell line models, for example, cell types not easily maintained in culture may be studied; gene expression during animal development may be studied; relevant mouse models of human disease may be developed; reporter mice may be crossed with mouse strains with other transgenes or mouse knockouts to evaluate the biological effects of these genes on telomerase RNA gene promoter activity; and compounds or molecules thought to modulate promoter activity can be tested *in vivo*, thus taking into account pharmacology of the test compounds. This may be important if modulators of promoter activity are to be used clinically.

In a further aspect, the present invention provides a method for identifying genes regulating telomerase RNA gene expression comprising the step of introducing reporter constructs stably into cell lines and single chromosomes into the cell lines via microcell-mediated chromosome transfer (England, N.L. et al Carcinogenesis Vol. 17(8) pp 1567-1575, 1996). This allows for fluctuations in reporter gene activity due to genes on the single chromosome to be monitored. This will identify genes carrying regulators of promoter expression and thus the signal transduction pathway controlling telomerase RNA

gene transcription. Gene products identified in this way will be considered as targets for inhibition and form part of the present invention.

5 The present invention provides methods for identifying genes regulating telomerase RNA gene expression comprising the steps of introducing reporter constructs stably into cell lines and expression libraries transfected into the reporter lines. The expression libraries may be made from cells known to repress or activate telomerase activity and
10 telomerase RNA expression. Fluctuations in reporter activity due to genes in the expression library may then be monitored, and the genes cloned. Gene products identified in this way will be considered as targets for inhibition and form part of the present invention.

15 In a further aspect, the present invention provides a method for testing candidate genes for promoter regulating activity, comprising the steps of transfecting candidate genes into cell lines containing the promoter/reporter constructs and monitoring fluctuations in reporter activity
20 due to said candidate genes.

The present invention extends to the use of the TR gene promoter for any of the methods given above. This includes the use of any promoter/reporter constructs, preferably those identified in Fig 4 and Fig. 5.

25 Aspects and embodiments of the present invention will now be further described by way of example only with reference to the accompanying drawings. Further aspects of the invention will be apparent to those or ordinary skilled in the art. All documents mentioned in this text are
30 incorporated herein by reference.

Brief description of the drawings.

Figure 1 shows the 1765bp genomic nucleotide sequence (SEQ I.D. No. 1) of human telomerase RNA gene (hTR) encompassing the gene promoter region. Transient expression of hTR-reporter gene constructs in HeLa and GM847 cells identified indicate that the elements responsible for promoter activity are contained in a 231bp region upstream of the transcriptional start site.

Figure 2 shows the 4044bp genomic nucleotide sequence (SEQ I.D. No. 2) of mouse telomerase RNA gene (*terc*) encompassing the gene promoter region. Transient expression of *terc*-reporter gene constructs in SWISS3T3 and A9 cells identified the elements responsible for promoter activity are contained in a 73bp region upstream of the transcriptional start site.

Figure 3 shows the restriction enzyme map of the genomic clones encompassing the human and mouse telomerase RNA genes. A) shows human telomerase RNA gene, hTR, genomic structure, 1765bp; B) shows mouse telomerase RNA gene *terc*, genomic structure. The transcribed regions of hTR and *terc* are depicted as black boxes within the central regions of the genomic sequences and the site of the template sequence within the telomerase RNA genes is indicated. The position of the CpG islands are shown as a box beneath the genomic sequence. Numbers in brackets refer to the nucleotide position within the sequence. The 3'-end of all the human promoter fragments is shown as hProm and fragments extend 5-prime to hProm867, hProm697, hProm341 and hProm111. The 3'-end of all the mouse promoter fragments is shown as mProm and fragments extend 5-prime to mProm628, mProm458, mProm418, mProm267,

mProm208, mProm136. The numbers after the prefix, hProm or mProm refer to the number of nucleotides of genomic sequence contained in the promoter fragment.

Figure 4 shows the nucleotide sequence of the human, (a), and mouse, (b), telomerase RNA gene 5'-flanking regions. Putative regulatory motifs are underlined. Arrows indicate the transcriptional start sites, (Blasco et al., 1995; Feng et al., 1995) and numbers to the left of each Figure refer to the number of bases upstream of the transcriptional start site. The template regions are in bold and underlined. Sequences contained in promoter constructs are shown by vertical lines and labelled, hProm or mProm, (see Figures 1 and 3). The regions containing elements responsible for minimum promoter activity are highlighted in bold, (see text for details). The run of CpA dinucleotide repeats in the mouse promoter is shown in bold and italic.

Figure 5 shows the detection of promoter activity in the 5'-flanking regions of human and mouse telomerase RNA genes. For each construct, the length of sequence upstream from the transcriptional start site is shown to the left and the luciferase activity to the right. a) Diagram comparing luciferase activity from human promoter constructs in GM847 and HeLa cells. Data for each construct is plotted as a percentage of the hProm505 luciferase activity as this construct consistently gave the highest activity in human cells. For each construct the mean and standard deviation for duplicate transfected wells is shown. b) Diagram comparing luciferase activity from mouse promoter constructs in SWISS3T3 and A9 cells. Data for each construct is plotted as a percentage of the

mProm458 luciferase activity as this construct consistently gave the highest activity in mouse cells. For each construct the mean and standard deviation for duplicate transfected wells is shown.

5 Figure 6 shows details of the oligonucleotide primers used for both the human and the mouse sequences. The positions of the primers can be seen on the maps provided in Figures 7 and 8.

10 Figure 7A shows a map of the 1765bp hTR construct with the position of the primers (see Fig. 6) shown. In each case the percentage homology of the primer is indicated.

 Figure 7B shows a map of the 1765bp hTR construct with the restriction sites marked.

15 Figure 8A shows a map of the 4044bp mouse *terc* construct with the position of the primers (see Fig. 6) shown. In each case the percentage homology of the primer is indicated.

 Figure 8B shows a map of the 4044bp mouse *terc* construct with the restriction sites marked.

20 Figure 9 shows the sequence of the TR gene promoter from Balb/c clones. The sequence analysis shows that the sequence is identical to that of the P1 sequence (Fig. 4b) apart from minor polymorphisms.

25 Figure 10 shows potential transcription factor binding sites within the hTR promoter.

30 Figure 11 shows the basal hTR promoter (-107 to + 69). This is a 176bp region termed sequence 2923 which contains several potential Sp1 transcription regulation sites, retinoblastoma control elements, (RCE) and the promoter CCAAT-box transcriptional regulator site. Sequence elements are defined in Figure 12.

Figure 12 shows oligonucleotides used in the cloning and mutagenesis of the hTR promoter region.

Figure 13 concerns mutagenesis of CCAAT-binding site. It shows a gel shift assay with the h10 sequence for the CCAAT binding site. h10 can bind to the Hela nuclear extract protein and can be competed by itself (h10). Mutation (h10m1) does not compete with oligo(h10) whereas h10m2 still competes. CCAAT site was mutated from CCAA to AGTC leading to loss of binding activity. This suggests that CCAAT site is a functional region for DNA-binding proteins.

Figure 14 shows nuclear factors binding to the Sp1 sites in the hTR promoter. Two separate DNA regions can be recognised by Hela protein extract and produce one Sp1(top) band-shift, two Sp3 (bottom) band-shift. This shows two Sp1 binding sites present in the proximal hTR promoter region.

Figure 15 shows site-directed mutagenesis of the Sp1-2 binding sites. It shows that specific Sp1 binding can be blocked by two bp nuclear changes in the Sp1-2(h41) sequence. 1: no competitor 2: competition by Sp1 consensus sequence 3: competition by Sp1-2 itself 4: no competition by mutation, h41m.

Figure 16 shows site-directed mutagenesis of the Sp1-1 binding region. It shows Sp1-1(h9) is critical region for hTR promoter. A series of mutations were introduced into h9 region by four bp change. Mutations are h9m1, h9m11, h9m2, h9m21 and h9m3. First lane no competitor. Second lane, competition by Sp1-1 itself(h9); fifth and six lanes show no competition by mutation m2 and m21; third, fourth and seventh lanes show competition by mutation 1, 11 and mutation 3. This suggests that mutation m2 and m21 base

pair region responsible for DNA-binding protein activity. Mutation of TATA-box still competes wild type DNA-binding.

Figure 17 concerns functional analysis of promoter mutants. It shows the scheme for introducing specific mutations into the hTR promoter. In brief, the mutant fragments were cloned into the luciferase basic vector, and the mutant confirmed by sequence. Mutant constructs were then transfected into 5637 cells to analysis luciferase activity. 5637 cells are a bladder carcinoma cell line which is p53 and pRb negative.

Figure 18 shows deletion analysis of the 5' flanking region of the hTR gene in Hela cells. The wild type promoter sequence used in this study is the hTR 2923 sequence shown in Figure 11. The promoter activity of this clone relative to the previously described promoter clones, (see Fig 5a), is shown in Figure 18.

Figure 19 concerns scanning mutation analysis of the hTR proximal promoter region (-107 to +69). It shows constructs with hTR promoter sequence element mutations. The parental wild type sequence is called 2923 and mutant constructs are shown below.

Figure 20 shows a diagrammatic version of the mutant promoter constructs described in Figure 19, which were used to assay for promoter activity.

Figure 21 concerns mutation analysis of the hTR promoter in 5637 cells. It shows the ability of the mutant promoters to drive gene expression in comparison to the wild type promoter sequence in cell lines. Left side shows the mutation constructs, right side shows luciferase activity relative to wild type(2923). Mutation Sp1-2 decrease the basal activity. Mutation Sp1-1 or both Sp-1

and Sp1-2 significantly increase the basal activity about 4 to 4.5 fold. Mutation CCAAT site abolished promoter activity completely. This result suggests that CCAAT-binding factor is essential element for basal hTR promoter activity. Two Sp1 elements involved in the hTR gene regulation. Sp1-1 is a site for negative regulation.

Figure 22 shows Sp1 activation of the hTR promoter in 5637 cells. Left side shows mutant constructs, right side shows related luciferase activity, right column shows fold induction by Sp1.

Figure 23 shows Sp3 repression of the hTR promoter in 5637 cells. Left side shows mutant constructs, right side shows related luciferase activity, right column show fold repression by Sp3.

Figure 24 concerns pRB interaction with Sp1 family and CCAAT factors. It shows activation of the hTR promoter in 5637 cells by the retinoblastoma gene product. Left side shows mutant constructs, right side shows related luciferase activity, right column shows fold induction by pRb.

Figure 25 shows a summary of the transcriptional regulation of the hTR gene promoter by Sp1, Sp3 and pRb. Two constructs were used, hTR 2923 and hProm867 as described in Figure 18. Basic refers to the promoter activity of the constructs in the absence of Sp1, Sp3 or pRb. As can be seen, SP1 and Rb stimulate hTR gene expression and Sp3 supresses hTR gene expression to background.

Figure 26 shows a map of the hTR suicide gene vector pGT52-codAuppTelo and the genetic components of the vector.

Figure 27 shows details of the two transcriptional

units of the hTR suicide gene vector pGT52-codAuppTelo. The pGT52-codAuppTelo vector consists of two transcriptional units. First Transcriptional Unit contains the following parts: hTR, telomerase RNA gene promoter; PLAP, Placental Alkaline Phosphatase reporter gene; Tk, HSV thymidine kinase conferring sensitivity to ganciclovir; Sh ble, Zeocin antibiotic resistance gene. Second Transcriptional Unit contains the following parts: hEF1-HTLV promoter, elongation factor 1a & part of HTLV promoter; codA::upp, bacterial cytosine deaminase & uracil phosphoribosyltransferase conferring sensitivity to the drugs 5FU & 5-fluorocytosine.

First Transcriptional Unit

hTR: telomerase RNA gene promoter
PLAP: Placental Alkaline Phosphatase reported gene
Tk: HSV thymidine kinase - sensitivity to ganciclovir
Sh ble: Zeocin antibiotic resistance gene

Second Transcriptional Unit

hEF1-HTLV prom: elongation factor 1a & part of HTLV promoter
codA::upp: bacterial cytosine deaminase & uracil phosphoribosyltransferase - sensitivity to 5FU & 5-fluorocytosine.

Detailed description

Preparation of tissue sections for in situ hybridisation

Formalin fixed paraffin embedded tissue blocks were obtained from Pathology Department files. Tissue sections were deparaffinised, rehydrated through graded concentrations of ethanol, (100%, 90%, 70%, 50%, 30% EtOH,

10 sec. each), rinsed in 0.85% sodium chloride for 5 minutes, followed by PBS for 5 minute. Sections were fixed in 4% paraformaldehyde/PBS for 20 minutes, rinsed in PBS, and treated with proteinase K (40 μ g/ml) in 50mM Tris-HCl pH 7.5, 5mM EDTA for 7.5 minutes at room temperature. After rinsing for 5 minutes in PBS, sections were post fixed in 4% paraformaldehyde/PBS for 5 minutes, rinsed in water, and acetylated in freshly prepared 0.25% acetic anhydride/0.1M triethanolamine for 10 minutes. The slides were rinsed in 0.85% saline, followed by PBS for 5 minutes each and dehydrated in gradually increasing concentrations of ethanol prior to hybridisation.

Probe preparation for in situ hybridisation

15 The riboprobe plasmid containing telomerase RNA sequences used for RNA in situ hybridisation is as previously described (Soder et al., 1997b). Control riboprobes were human histone H3 and glyceraldehyde 3-phosphate dehydrogenase (GAPDH), (Ambion, Texas). The probes were labelled with (³⁵S)-UTP using a RNA labelling kit (Amersham, UK). Transcripts were purified using a Sephadex G-50 column (Pharmacia), phenol/chloroform extracted and precipitated in ethanol. The probes were resuspended in 50 mM dithiothreitol. Northern blot analysis of normal human tissue confirmed the specificity and sensitivity of the hTR probe to detect hTR expression in normal testis, (data not shown).

Hybridisation and washing procedures

30 Sections were hybridised overnight at 52°C in 60% formamide, 0.3M NaCl, 10mM Tris-HCl (pH 7.5), 5mM EDTA, 10%

dextran sulphate, 1xDenhardt's (0.02% BSA, 0.02% Ficoll, 0.02% polyvinylpyrrolidone), 0.5mg/ml yeast tRNA, 50mM DTT (freshly added), and 50 000 cpm/ul ³⁵S-labelled cRNA probe. The tissue was washed stringently at 50°C in 5x SSC, 0.1% β-mercaptoethanol for 25 minutes, at 65°C in 50% formamide, 2x SSC, 1% β-mercaptoethanol for 25 minutes, and washed twice at 37°C in 0.5M NaCl, 10mM Tris-HCl pH 7.5, 5mM EDTA for 15 minutes before treatment with 20μg/ml RNaseA at 37°C for 30 minutes. RNase A only digests single stranded RNA. This removes single stranded, and therefore unhybridised probe, leaving the RNA:RNA duplexes intact. Thus this step helps reduce background probe signal. Following washes in 50% formamide, 2xSSC, 1% β-mercaptoethanol at 65°C for 20 minutes, and twice in 2x SSC at 50°C for 15, the slides were dehydrated and dipped in 0.1% gelatine/0.01% chromealum, and then in Hypercoat Nuclear LM-1 emulsion (Amersham) and exposed for 2 weeks in light tight boxes with desiccant at 4°C. The microautradiographs were developed in 20% Phenisol for 2.5 minutes, washed in 1% acetic and water each for 30 seconds, fixed in 30% sodium thiosulphate for 5 minutes, rinsed in water 30 minutes, and counter-stained with haematoxylin.

Cloning of sequences encompassing the human and mouse telomerase RNA genes

The present inventors have previously reported the identification of genomic clones in P1 vectors containing hTR and *terc* transcribed sequences, (Soder et al., 1997b; Soder et al., 1997c). The human P1 clone, 9913, is derived from a human foreskin fibroblast P1 library and the mouse P1 clone, 11792, is derived from a mouse C127 fibroblast P1

library. Briefly, in order to subclone the promoter regions, the P1 clones were digested with *EcoRI* and *HindIII* and ligated into pBluescript. Colonies containing telomerase RNA gene sequences were identified by hybridisation with PCR generated probes as previously described, (Soder et al., 1997b; Soder et al., 1997c). Plasmid DNA was prepared from positively hybridising colonies, and inserts sequenced on both strands by dideoxy chain termination using the ABI PRISM dye terminator cycle sequencing kit (PE Applied Biosystems, Warrington, UK) and 25ng oligonucleotide primers. Dye labelled products were resolved and detected using the Applied Biosystems DNA sequencer ABI373. Sequence was analysed using the Sequencing Analysis program v3.0. Homology searches carried out using BLAST, (Basic Local Alignment Search Tool), National Centre for Biotechnology Information, (NCBI): <http://www.ncbi.nlm.nih.gov/>. Sequence was analysed for potential transcription factor binding sites by TESS: Transcription Element Search Software on the WWW, Jonathan Schug and G. Christian Overton, Technical Report CBIL-TR-1997-1001-v0.0, of the Computational Biology and Informatics Laboratory, School of Medicine, University of Pennsylvania, 1997, <http://agave.humgen.upenn.edu/tess/index.html>. Identification of CpG islands was carried out using GRAIL: Gene Recognition and Assembly Internet Link, <http://compbio.ornl.gov/Grail-1.3/>. The full sequences have been submitted to GenBank.

Construction of luciferase reporter gene constructs

The structures of the telomerase RNA gene-luciferase constructs used in the study are shown in Figures 5 and 4.

Promoter-luciferase constructs were made by inserting PCR products into pGL3-Basic, (Promega). Orientation and sequence of each insert was checked by sequencing. Details of the primers are given in Figure 6.

5

Transfection and luciferase assays

All transfections were carried out in duplicate in 6-well plates, (35mm diameter). Cells were seeded at 6×10^4 cell per well and cultured overnight. Transfection was carried out using SuperFect Transfection Reagent, (Qiagen), according to the manufacturers instructions. Cells were exposed to the transfection mix for three hours and harvested for analysis after 48 hours. Equivalent amounts of cellular protein as determined by Bio-Rad assay, (BioRad), were used in the luciferase assay. Luciferase assays were performed according to the manufacturers protocols, (Promega). To ensure reproducibility in the assays, particular care was taken over the following: DNA used for transfection was quantified by spectrophotometry and direct visualisation by gel electrophoreses. All transfections were carried out in duplicate wells and this was found to be a good measure of the reproducibility of transfection. In each experiment, all deletion constructs were analysed together with both the basic cloning vector, pGL3-Basic and the positive control vector, pGL3-Control, which contains SV40 promoter and enhancer sequences. Each extract was measured for luciferase activity at least twice. All transfections were carried out at least three times. Initial transfection conditions were determined by using promoter fragments liked to a green fluorescent protein reporter gene, (Clontech), as this allowed direct

30

visualisation of promoter activity in live cells, (data not shown). The present inventors found it important to transfect and analyse the cells at sub-confluence and that it was important to avoid harsh transfection protocols such as electroporation, resulting in poor cell viability.

Tumour specific regulation of telomerase RNA gene expression visualized by in situ hybridization.

The patterns of hTR expression were examined in epithelial cancer of lung, ovary, breast and cervix, (Table 1). Twenty six percent of non-small cell lung cancers, (NSCLC), were hTR positive. However, the NSCLC group consists of squamous, adenocarcinoma and large-cell anaplastic variants. Interestingly however, expression was almost exclusively limited to the squamous variants, (p=0.006), with 41% of squamous NSCLC expressing hTR but only 8% of adenocarcinoma and large-cell anaplastic NSCLC expressing hTR. This data suggests that hTR may be differentially regulated during the oncogenesis of squamous and non-squamous NSCLC. Indeed, the low frequency of detectable hTR expression in adenocarcinoma of the lung was also observed in adenocarcinomas of ovary and breast, (Table 1). In addition, metastatic carcinoma in hilar lymph nodes of 19 of the NSCLC cases were available for a comparative study with the paired primary carcinomas. All 6 cases which expressed hTR in the primary carcinoma retained expression in the metastasis and all 13 primary carcinomas which lacked detectable hTR expression remained negative in the metastasis, (Table 1). Thus, expression levels appear stable between primary and metastatic carcinomas and expression of hTR is not associated with

metastasis of pulmonary carcinomas.

Cancer of the uterine cervix is a heterogeneous group of lesions, which like NSCLC can be subdivided into squamous and adenocarcinoma (Benda, 1994). The present
5 inventors studied 87 cervical lesions for hTR expression, (Table 1). hTR expression was detected in 44% of the cervical carcinomas, however in contrast to NSCLC, there was no significant difference in frequency of expression between invasive squamous carcinoma, (44%), and invasive
10 adenocarcinoma, (32%). The data for adenocarcinoma of the cervix also contrast those for invasive adenocarcinoma of the breast, (13%), and ovary, (17%), (Table 1), and suggest that regulation of hTR expression may be different for cervical cancer and therefore relate to the aetiology of the disease (Benda, 1994; Klingelhutz et al., 1996).
15 Interestingly, hTR expression was readily detected in preinvasive cervical cancer, (40%, see Table 1), and there was no significant difference in frequency between invasive and preinvasive lesions. In addition, the case of glandular
20 intraepithelial neoplasia of the cervix has heterogeneous expression of hTR (data not shown), thus allowing the evolution of hTR expressing cells to be followed in their histological context.

The primitive germ cells of the male are found in the
25 seminiferous tubules and the present inventors examined 22 sections from normal testis and uninvolved tubules from testicular cancer patients to establish the pattern of hTR expression in normal seminiferous tubules, (data not shown). Of the 22 sections, 21 showed hTR expression in
30 the primitive germ cells located in the basal layers of the seminiferous epithelium, (Table 1). The intimate

relationship between germ cells and the supportive Sertoli cells, means that expression of hTR in the Sertoli cells cannot be excluded. Mature germ cells (spermatids and spermatazoa) when present, did not express hTR. Thus, the

5 *in situ* assay can detect normal levels of hTR expression in primitive germ line stem cells and the distribution of hTR expression in the testis is consistent with its proposed role in the maintenance of telomere length in the germ line. A series of 22 testicular germ cell tumours were

10 also analysed for hTR expression. As shown in Table 1, 73% of testicular germ cell tumours were positive for hTR expression, and there was no significant difference between teratomas and seminomas of the testis. Interestingly, within hTR-positive teratomas, mature tissues never had

15 detectable hTR expression whatever their differentiation. Similarly, 15 benign ovarian teratomas composed of fully mature differentiated tissues had no detectable hTR expression, (Table 1). Thus, the differentiated teratomas may recapitulate the down regulation of hTR during early

20 embryonic or foetal development, (Wright et al., 1996). In five of the testicular seminomas there was no detectable hTR expression, (Table 1), but in each seminoma case, germ cells within adjacent normal seminiferous tubules expressed hTR, suggesting either that hTR expression has been

25 repressed during oncogenesis, or that seminomas without detectable levels of hTR arise from germ cells with low or no hTR expression.

Table 1. Frequency of hTR expression.

30

Tumour	Tissue histology	Frequency	%

	Testicular Germ Cell Tumours	<i>normal testicular germ cells</i>	21/22	95
		^a teratoma	7/8	
5		seminoma	8/13	
		intratubular	1/1	
		Total	16/22	73
10	Ovarian Germ Cell Tumours	^b <i>benign ovarian teratomas</i>	0/15	0
	Non-small Cell Lung Cancer	squamous	13/32	41
15		adenocarcinoma(18)/large cell(7)	2/25	8
		^c Total	15/57	26
	Epithelial Ovarian Cancer	^d <i>adenocarcinoma</i>	6/34	17
20				
	Breast Cancer	^e invasive adenocarcinoma	7/54	13
		ductal carcinoma in situ, (DCIS)	0/19	0
		^f DCIS, recurrent	1/4	
		Phylloides	0/6	
25		Total	8/83	10
	Cervical Cancer	invasive squamous	15/34	44
		invasive adenocarcinoma	7/22	32
		^g invasive adenosquamous	7/11	
30		Invasive Total	29/67	43

	squamous intraepithelial neoplasia	4/5	
	glandular intraepithelial neoplasia	2/10	
	^b mixed squamous/glandular intraepithelial neoplasia	3/5	
	Preinvasive Total	8/20	40
5			
	^c All cervix samples	38/87	44
	Metastasis:	positive in primary & metastasis	6/6
10	Non-small Cell Lung Cancer	negative in primary & metastasis	13/13

Table 1. Frequency of hTR expression. ^aThe single teratoma without detectable hTR expression was of intermediate maturity and also expressed GAPDH, (data not shown), and therefore the lack of expression is not due to loss of RNA from the sample. ^bTo test for RNA integrity in the ovarian teratomas, a group of 6 were analysed for GAPDH expression and all were found to be positive. ^cA group of 33 NSCLC were tested for histone H3 expression, 22 of which did not have detectable levels of hTR expression and all 33 were positive. An additional 4 hTR negative NSCLC samples tested for GAPDH expression were positive. ^dTwo ovarian cancers, (1 hTR negative), were tested for histone H3 expression and both were positive. Three ovarian cancers were tested for GAPDH expression and all were positive. ^eA group of 10 invasive breast cancers, (7 hTR negative), were tested for histone H3 expression and all were positive. In addition, 11 breast cancers lacking detectable levels of hTR expression were tested and were positive for GAPDH expression. ^fFour specimens of DCIS were obtained from patients who had in the previous 5 years been diagnosed and treated for the presence of preinvasive breast cancer, (DCIS). The primary biopsies for all 4 were lacking detectable levels of hTR expression, however one recurrent DCIS biopsy had detectable hTR expression. ^gAdenosquamous cancers of the cervix. Where samples

were positive for hTR expression, all elements were positive. ^bPreinvasive cervical cancer. In lesions with mixed glandular and squamous elements, where samples were positive for hTR expression, all elements were positive. Two cervical cancers, (1 lacking detectable levels of hTR expression), were tested for histone H3 expression and both were positive. Three cervical cancers were tested for GAPDH expression and all were positive. In total, 28 samples, (25 lacking detectable levels of hTR expression), were tested and were positive for GAPDH expression and 47 samples, (31 lacking detectable levels of hTR expression), were tested and were positive for histone H3 expression. Visual inspection suggested that histone H3-positive cells were present in comparable numbers in the carcinomas lacking detectable levels of hTR expression and in the hTR positive carcinomas. The same areas of tissue were examined for both hTR and histone H3 expression, thus, the lack of detectable hTR expression is unlikely to be due simply to the presence of quiescent cells in the tumours, (Holt et al., 1996b).

Cloning of genomic sequences encompassing the human and mouse telomerase RNA genes.

In order to obtain sequences flanking the genes, the P1 genomic clones were digested with *EcoRI* and *HindIII*, subcloned into the plasmid vector, pBluescript, and colonies containing hTR or *terc* sequences identified by hybridisation to PCR generated probes specific for the genes. A 1.3kb genomic clone encompassing hTR was isolated as was a 4kb genomic clone encompassing *terc*, (Figure 3).

A BLAST search using the 1.3kb human sequence identified three high-scoring segment pairs: HSU85256, HSU86046 and MMU33831. HSU85256, (598bp of sequence), and HSU86046, (545bp of sequence), are published sequences for the transcribed region of the human telomerase RNA gene and confirmed that we had cloned genomic sequences encompassing hTR (Bryan et al., 1997; Feng et al., 1995). MMU33831 is the sequence of the transcribed region of the mouse telomerase RNA gene which has previously been shown to have homology to the human gene (Blasco et al., 1995).

A BLAST search using the 4kb mouse sequence identified both the published human gene sequences, (HSU85256, HSU86046), and the published sequence for the transcribed region of the mouse gene, MMU33831, (591bp of sequence) (Blasco et al., 1995; Bryan et al., 1997; Feng et al., 1995). In order to confirm that the genomic sequence obtained from the P1 subclone was genuine, the present inventors cloned 5'-flanking sequences using genomic DNA from Balb/c mice in PCR reactions. Sequence analysis of the Balb/c clones were identical to the P1 sequence except for minor polymorphism's, (Fig. 9). A schematic representation of the 4kb of sequence information encompassing the mouse telomerase RNA gene is shown in Figure 3.

15

Analysis of nucleotide sequence encompassing the human and mouse telomerase RNA genes.

To investigate the relationship between the human and mouse genomic clones, sequence comparisons were carried out. The transcribed regions of the two genes showed 67% identity in keeping with the published estimate, (Feng et al., 1995). However, no significant sequence identity could be identified in either the 5'- or 3'-regions flanking the transcribed sequences.

25

Both the human and mouse sequences were analysed for CpG islands by GRAIL. CpG islands were defined as regions larger than 200bp, with an average GC content greater than 50% and the ratio of observed versus expected CpGs greater than 0.6 (Gardiner-Garden & Frommer, 1987). Interestingly, both the human and mouse genes lie within CpG islands, (see Figure 3). The human gene is covered by a CpG island 733bp

30

in length, with a GC content of 66% and a ratio of observed versus expected CpGs of 0.89. The mouse gene is covered by a CpG island of 659bp in length, with a GC content of 64% and a ratio of observed versus expected CpGs of 0.81.

5 The 5'-flanking regions of the human and mouse telomerase RNA genes were also analysed for potential transcription factor recognition sites. As shown in Figure 4, a number of potential binding sites can be identified, including consensus sequences for
10 glucocorticoid/progesterone/androgen receptor binding, AP1 and Ets family members. CCAAT box's are found in both genes close to the published transcriptional start sites (Blasco et al., 1995; Feng et al., 1995), however, there is no obvious TATA box in the mouse gene with the human gene
15 TATA box consensus sequence being in the reverse orientation. The mouse promoter region also contains a run of CpA dinucleotide repeats which may be of use in developing microsatellite genetic markers for this gene.

20 Transfection assays detect promoter activity in the 5'-flanking regions of the human and mouse telomerase RNA genes.

To identify whether the 5'-flanking DNA of the telomerase RNA genes exhibited promoter activity, sequences
25 were fused to a firefly luciferase reporter gene, (pGL3-Basic). The transcriptional start sites for both the human and mouse telomerase RNA genes have been established (Blasco et al., 1995; Feng et al., 1995). Various promoter constructs containing the transcriptional start site were
30 therefore generated, (see Figures 4, 5).

Human promoter constructs containing truncated

portions of the 5'-flanking region were transiently transfected into HeLa and GM847 cells, (Figure 5a). HeLa is a telomerase positive cervical carcinoma cell line, GM847 is a SV40-immortalised skin fibroblast cell line which expresses the telomerase RNA component but is telomerase-negative, (Bryan et al., 1997). As shown on Figure 5a, promoter activity was observed in both cell lines with fragments containing 341bp or more, (from position -272, see Figure 4a, 5a). The highest luciferase activity was observed with construct hProm505 which contains a 505bp fragment, (position -463, see Figure 4). Construct, hProm111, which contains only 111bp of 5'-flanking sequence, (position -42, see Figure 4a, 5a), produced a dramatically reduced level of luciferase activity, (Figure 5a). Thus, a minimal promoter sequence can be defined as extending 272bp upstream of the transcription start site, and that elements responsible for promoter activity must be contained in a 231bp region between -272bp and -42bp, (Figures 4a and 5a).

Mouse promoter constructs containing various truncated portions of the 5'-flanking region were transiently transfected into SWISS3T3 and A9 cells, (Figure 5b). SWISS3T3 cells are an embryo derived line and A9 cells are of areolar and adipose origin. Both cell lines are telomerase positive. As shown on Figure 5b, promoter activity was observed in both cell lines with fragments containing 208bp or more, (from position -94, see Figure 4b). Construct mProm136, which contains only 136bp of 5'-flanking sequence, (position -22, see Figure 4b), produced dramatically reduced levels of luciferase activity, (Figure 4b). Thus, a minimal promoter sequence can be defined as

extending 94bp upstream of the transcription start site, and that elements responsible for promoter activity must be contained in a 73bp region between -94bp and -22bp, (Figure 4b and 5b).

5 Transfection of the human promoter construct hProm867 into mouse cells gave very strong promoter activity, with up to twice that of the strongest mouse construct, and transfection of the mouse promoter construct, mProm628 into human cells also showed luciferase activity at around 25%
10 of the strongest human construct, (data not shown).

Identification of Transcription Factor Binding Sites in the hTR Promoter by Gel Shift assays

In order to identify the key sequence elements
15 involved in transcription factor binding, protein extracts from cell lines were assayed for their ability to detect specific DNA sequence elements within the hTR promoter by gel shift assays. Fluorescence assays may also be used and detected by microscopy, microplate reader or cytometry.
20 Such elements are then considered candidates for regulating hTR *in vivo*. These studies concentrated on the 176bp region of the hTR promoter shown in Figure 11 as part of the sequence shown in Figures 1 and 4a. This 176bp region is termed sequence 2923 and contains several potential Sp1
25 transcription regulation sites, retinoblastoma control elements, (RCE) and the promoter CCAAT-box transcriptional regulator site, (see Figure 11). The oligonucleotide sequence elements used in the analysis of transcriptional regulation are shown in Figure 12.

30

Identification of protein interaction with the CCAAT

sequence element.

A CCAAT box element is found upstream of the transcriptional start site for hTR, (see Figure 11). In order to identify whether cellular proteins interact with this site in the hTR promoter, protein extract from the HeLa cell line was used in an electrophoretic mobility shift assay, (EMSA) with oligonucleotide sequence elements corresponding to the normal wild type hTR sequence termed h10 in Figure 12 & 13, (see Figures 11, 12 & 13). Figure 13 shows that the h10 oligonucleotide sequence containing the hTR CCAAT-box can bind proteins in a specific fashion as competition for binding with the h10 oligonucleotide sequence itself abolishes the complex as does an oligonucleotide with mutations outside the CCAAT region, (see sequence h10m2, Figures 12 & 13). Addition of oligonucleotide sequences with mutations introduced into the CCAAT site cannot compete for protein binding, (sequence h10m1 in Figure 12 & 13). This suggests that the CCAAT-box sequence element is a functional region for DNA binding proteins and may regulate the hTR promoter.

Identification of Two Sp1 sequence elements in the hTR promoter.

Two Sp1 sequence elements are found upstream of the hTR transcriptional start site, (Figure 11). In order to identify whether cellular proteins interact with these sites in the hTR promoter, protein extract from the HeLa cell line was used in an electrophoretic mobility shift assay, (EMSA) with oligonucleotide sequence elements corresponding to the normal wild type hTR sequences termed h4, (Sp1-2, see Figures 11 & 12), and h9, (Sp1-1, see

Figures 11 & 12). As can be seen from Figure 14, 15, 16, hTR promoter sequence containing Sp1-1 and Sp1-2 sequence elements form DNA/protein complexes characteristic of binding of Sp1 and Sp3 transcription factors. This suggests that the hTR promoter may be regulated by the Sp family of transcription factors.

Functional Analysis of hTR Promoter Sequence Elements in Cell Lines

The gel shift assays discussed above identified a number of sequence elements which form specific DNA/protein complexes which may regulate the activity of the hTR promoter. In order to confirm these sequence elements function in the regulation of the hTR promoter *in vivo*, mutations were introduced into these sites within the context of the basal promoter region of the hTR gene. The ability of these mutations to drive the hTR promoter in comparison to the wild type promoter sequence was then analysed. The overall scheme of this approach is shown in Figure 17.

The wild type promoter sequence used in this study is the hTR 2923 sequence shown in Figure 11. The promoter activity of this clone relative to the previously described promoter clones, (see Fig 5a), is shown in Figure 18. In order to analyse the function of hTR promoter sequence elements in cell lines, constructs with hTR promoter sequence element mutations were made and these are shown in Figures 19 & 20. The ability of the mutant promoters to drive gene expression was assayed in comparison to the wild type promoter sequence in cell lines as shown in Figure 21. Figure 21 shows firstly that mutation of the Sp1-2 sequence

element reduces promoter activity, (construct 29m223),
sugesting this element is the site of action for an
activator of the hTR promoter; secondly that mutation of the
Sp1-1 sequence element significantly increases basal
5 promoter activity by 4-4.5 fold, (construct 92 and 29m292),
sugesting this element is the site of action for an
inhibitor of hTR promoter activity; and thirdly that
mutation or deletion of the CCAAT sequence element
abolishes hTR promoter activity, (costruct 1011 & 26n23),
10 suggesting this element is the site of action for an
activator of hTR promoter activity.

Thus the inventor has identified key regulatory
sequence elements involved in hTR promoter activation and
repression.

15

Identification of Genes Modulating hTR Promoter Activity

In order to identify genes regulating telomerase RNA
gene expression, candidate genes are transfected into cell
lines containing the hTR promoter/reporter constructs.
20 Fluctuations in reporter activity due to candidate genes
can are monitored and conclusive evidence presented for
their activity.

The analysis of hTR sequence elements presented above
sugests the involvement of the Sp family of transcription
25 factors in regulation of the hTR promoter. Therefore, the
inventor co-transfected into cell lines the luciferase
reporter gene under control of the hTR promoter together
with expression vectors for the following three
transcriptional regulators: Sp1, Sp3, and pRb.

30 The transcriptional regulators Sp1 and Sp3 both
recognise the same DNA sequence, the so called Sp1 site.

Transcriptional regulation by the Sp family of proteins is complex, but in general Sp1 activates genes whereas Sp3 can repress promoter activity. The retinoblastoma gene product, pRb, is classically thought of as a tumour suppressor gene and has numerous roles as a modulator of gene transcription and cell cycle regulator. The Rb gene was the first tumour suppressor gene to be cloned. pRb has roles in: mediating transcriptional repression of RNA polymerase II, (Pol II), transcribed promoters; mediating transcriptional repression of RNA polymerase III, (Pol III), transcribed promoters; mediating transcriptional activation of RNA polymerase II, (Pol II), dependant promoters. This last role in mediating transcriptional activation is possibly the least well understood and studied. However, many important genes are activated by pRb, such as, TGF-B2, Il-6, the retinoblastoma gene itself, and cyclin D1, (Herwig & Strauss, 1997; Lania et al., 1997; Sellers & Kaelin, 1996; Tenen et al., 1997).

The Sp1 Transcription Factor Activates the hTR promoter.

The ability of the Sp1 transcription factor to modulate the hTR promoter was assessed by co-transfecting the hTR promoter constructs shown in Figure 22, with an Sp1 expression vector. From Figure 22 it can be seen that Sp1 is an activator of the hTR promoter.

The Sp3 Transcription Factor Represses the hTR Promoter.

The ability of the Sp3 transcription factor to modulate the hTR promoter was assessed by co-transfecting the hTR promoter constructs shown in Figure 23, with an Sp3 expression vector. From Figure 23 it can be seen that Sp3 is an repressor of the hTR promoter.

The Retinoblastoma Gene Product, pRb, Activates the hTR Promoter

The ability of the pRb to modulate the hTR promoter was assessed by co-transfecting the hTR promoter constructs shown in Figure 24, with an pRb expression vector. From Figure 24 it can be seen that pRb is an activator of the hTR promoter. For a summary of the above data see Figure 25.

10 Genetically directed enzyme prodrug therapy (GDEPT).

The term "GDEPT" is used to include both viral and non-viral delivery systems. Examples of suitable vector systems include vectors based on the Molony murine leukaemia virus are known (Ram. Z et al, Cancer Research (1993) 53. 83-88; Dalton and Treisman, Cell (1992) 68. 597-612). These vectors contain the Murine Leukaemia virus (MLV) enhancer system cloned upstream at the β -globin minimal promoter. This vector further contains a polylinker to facilitate cloning, followed by the β -globin 3'-untranslated region and polyadenylation sites.

Suitable viral vectors further include those which are based upon a retrovirus. Such vectors are widely available in the art, Culver et al (Science (1992) 256. 1550-1552) also describe the use of retroviral vectors in GDEPT. Such vectors or vectors derived from such vectors may also be used. Other retroviruses may also be used to make vectors suitable for use in the present invention. Such retroviruses include rous sarcoma virus (RSV). The promoters from such viruses may be used in vectors in a manner analogous to that described for MLV.

In general the vector may be any RNA or DNA vector

used in VDEPT or GDEPT therapies.

The enzyme may be any enzyme which is not normally expressed in the surface of a cell, nor released into circulation, particularly a mammalian (especially human) cell, and which is capable of modifying or killing the target cell e.g. a cancer cell.

The enzyme may be linked to a signal sequence which directs the enzyme to the surface of a mammalian cell. This will usually be a mammalian signal sequence or a derivative thereof which retains the ability to direct the enzyme to the cell surface. This will be needed unless the enzyme has an endogenous signal sequence which does this. Even if the enzyme does have such a signal sequence, it can be replaced by another signal sequence where this is desirable or appropriate. Suitable signal sequences include those found in transmembrane receptor tyrosine kinases such as the c-erbB2 (HER2/neu) signal sequence or variants thereof which retain the ability to direct expression of the enzyme at the cell surface. The c-erbB2 signal sequence can be obtained by reference to Coussens et al (1985) Science. 230. 1132-1139.

The enzyme may be expressed at the surface of a cell. This means that it will be expressed in such a fashion as to have the enzyme exposed outside the cell so that it may interact with the prodrug, but will still be attached to the plasma membrane by virtue of a suitable plasma membrane anchor. A suitable anchor will be a polypeptide anchor which is expressed by the vector. For example, the enzyme which is linked to a sequence which is a transmembrane region which anchors the enzyme in the membrane of the cell. Such a transmembrane region can be derived from

transmembrane receptor kinases, such as cerbB2, EGF receptors and CSF-1 receptors.

Vectors encoding the enzyme, together with, when required, a signal sequence and/or transmembrane region may be made using recombinant DNA techniques known per se in the art. The sequences encoding the enzyme, signal sequence and transmembrane regions may be constructed by splicing synthetic or recombinant nucleic acid sequences together, or by modifying existing sequences by techniques such as site directed mutagenesis. Reference may be made to "molecular cloning" by Sambrook et al (1989) Cold Spring Harbor) for discussion for standard recombinant DNA techniques.

The enzyme will be expressed in the vector using a promoter sequence or functional part thereof according to the present invention capable of activated in the cell to which the vector is targeted. The promoter sequence according to the present invention will be operably linked to the enzyme and its associated sequences. The promoter sequences according to the present invention may be modified by techniques known in the art. Certain regions of the promoter sequence must be retained to ensure cell, e.g. tumour cell, specificity whereas other regions may be modified or deleted without significant loss of specificity (see Figure 5). The degree of regulation of such candidate promoter regions as described herein, can be tested and assessed by techniques known to those skilled in the art.

"Operably linked" refers to a juxtaposition wherein the promoter sequence and the enzyme-coding sequence are in a relationship permitting the coding sequence to be expressed under the control of the promoter sequence. Thus,

there may be elements such as 5' non-coding sequence between the promoter sequence and coding sequence which is not native to either the promoter or nor the coding sequence. Such sequences can be included in the vector if
5 they do not impair the correct control of the coding sequence by the promoter.

The vectors for use in GDEPT or VDEPT systems comprising the promoter sequence or functional fragment thereof in accordance with the present invention can be
10 used in a method of treatment of the human or animal body. Such treatment includes a method of treating the growth of neoplastic cells which comprises administering to a patient in need of treatment the GDEPT or VDEPT systems. It is also possible that the vectors may be used to treat cells which
15 are diseased through infection of the human or animal body by bacteria, viruses or parasites.

For use of the vectors in therapy, the vectors will usually be packaged into viral particles and the particles delivered to the site of the tumour, as described in for
20 example Ram et al (ibid).

The viral particles maybe modified to include an antibody, fragment thereof (including a single chain) or tumour - directed ligand to enhance targetting of the tumour. Alternatively the vectors may be packaged into
25 liposomes. The liposomes may be targeted to a particular tumour. This can be achieved by attaching a tumour-directed antibody to the liposome. Viral particles may also be incorporated into liposomes. The particles may be delivered to the tumour by any suitable means at the
30 disposal of the physician. Preferably, the viral particles will be capable of selectively infecting the tumour cells.

By "selectively infecting" it is meant that the viral particles will primarily infect tumour cells and that the proportion of non-tumour cells infected is such that the damage to non-tumour cells by administration of a prodrug will be acceptably low, given the nature of the disease being treated. Ultimately, this will be determined by the physician.

One suitable route of administration is by injection of the particles in a sterile solution. While it is possible for the prodrugs to be administered alone it is preferably to present them as pharmaceutical formulations. The formulations comprise a prodrug, together with one or more acceptable carriers thereof and optionally other therapeutic ingredients. The carrier or carriers must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipients thereof, for example, liposomes. Suitable liposomes include, for example, those comprising the positively charged lipid (N[1-(2,3-dioleyloxy)propyl]-N,N,N-triethylammonium (DOTMA), those comprising dioleoylphosphatidylethanolamine (DOPE), and those comprising 3β [N-(n'-N'-dimethylaminoethane)-carbamoyl]cholesterol (DC-Chol).

Viruses, for example isolated from packaging cell lines may also be administered by regional perfusion or direct intratumoral direction, or direct injection into a body cavity (intracavitary administration), for example by intra-peritoneum injection.

Formulations suitable for parenteral or intramuscular administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants,

5 buffers, bacteriostats, bactericidal antibiotics and
solutes which render the formulation isotonic with the
blood of the intended recipient; and aqueous and non-
aqueous sterile suspensions which may include suspending
10 agents and thickening agents, and liposomes or other
microparticulate systems which are designed to target the
compound to blood components or one or more organs. The
formulations may be presented in unit-dose or multi-dose
containers, for example sealed ampoules and vials, and may
15 be stored in a freeze-dried (lyophilized) condition
requiring only the addition of the sterile liquid carrier,
for example water, for injections, immediately prior to
use. Injection solutions and suspensions may be prepared
extemporaneously from sterile powders, granules and tablets
of the kind previously described.

It should be understood that in addition to the
ingredients particularly mentioned above the formulations
may include other agents conventional in the art having
regard to the type of formulation in question. Of the
20 possible formulations, sterile pyrogen-free aqueous and
non-aqueous solutions are preferred.

The doses may be administered sequentially, eg. at
daily, weekly or monthly intervals, or in response to a
specific need of the patient. Preferred routes of
25 administration are oral delivery and injection, typically
parenteral or intramuscular injection or intratumoural
injection.

In using the system of the present invention the
prodrug will usually be administered following
30 administration of the vector incorporating the promoter of
the present invention and encoding an enzyme. Typically,

the vector will be administered to the patient and then the uptake of the vector by transfected or infected (in the case of viral vectors) cells monitored, for example by recovery and analysis of a biopsy sample of targeted tissue.

The exact dosage regime will, of course, need to be determined by individual clinicians for individual patients and this, in turn, will be controlled by the exact nature of the prodrug and the cytotoxic agents to be released from the prodrug but some general guidance can be given. Chemotherapy of this type will normally involve parenteral administration of both the prodrug and modified virus and administration by the intravenous route is frequently found to be the most practical. For glioblastoma the route is often intratumoural. A typical dosage range of prodrug generally will be in the range of from about 1 to 150 mg per kg per patient per day, which may be administered in single or multiple doses. Preferably the dose range will be in the range from about 10 to 75, e.g. from about 10 to 40, mg per kg per patient per day. Other doses may be used according to the condition of the patient and other factors at the discretion of the physician.

Tumours which may be treated using the system of the present invention include any tumours capable or being treated by a GDEPT or VDEPT system and thus are not limited to any one particular class of tumours. Particularly suitable tumour types include breast, colorectal and ovarian tumours, as well as pancreatic, melanoma, glioblastoma, hepatoma, small cell lung, non-small cell lung, muscle and prostate tumours.

The system of the invention may also be used to treat

infections diseases, for example, and any other condition which requires eradication of a population of cells.

It will be understood that where treatment of tumours is concerned, treatment includes any measure taken by the physician to alleviate the effect of the tumour on a patient. Thus, although complete remission of the tumour is a desirable goal, effective treatment will also include any measures capable of achieving partial remission of the tumour as well as a slowing down in the rate of growth of a tumour including metastases. Such measures can be effective in prolonging and/or enhancing the quality of life and relieving the symptoms of the disease.

Tumour specific gene expression for genetic therapy via telomerase RNA gene promoters.

The present studies point to examples of clear differentials in hTR expression between cancerous and adjacent normal tissue which support the application of effective telomerase-based therapy. Indeed, the presence of high levels of hTR expression in specific cancers suggest the TR promoter may be an interesting focus for genetic therapies designed to target therapeutic genes to tumours, via tumour specific gene expression. Therefore, the present inventor provide use of the TR promoters to drive expression of enzyme-prodrug activation systems such as viral thymidine kinase and ganciclovir, although many other systems may also be used (see above). Targeted gene expression via the telomerase RNA gene promoter may also be used in gene replacement strategies for cancer therapy.

To this end, the present invention provides a first generation suicide gene therapy vector using the hTR

promoter sequence, namely pGT52-codAuppTelo, (Figures 26 & 27). The hTR promoter sequences used to generate pGT52-codAuppTelo are hProm867 as shown in Figure 4a. These sequences were inserted into the commercially available gene therapy vector pGT62-codAupp, (Invivogen, San Diego, CA, USA), in place of the CMV promoter sequences.

The pGT52-codAuppTelo vector consists of two transcriptional units. First Transcriptional Unit contains the following parts: hTR, telomerase RNA gene promoter; PLAP, Placental Alkaline Phosphatase reporter gene; Tk, HSV thymidine kinase conferring sensitivity to ganciclovir; Sh ble (*Streptoalloteichus hindustanus* bleomycin gene), Zeocin antibiotic resistance gene. Second Transcriptional Unit contains the following parts: hEF1-HTLV promoter, elongation factor 1a and part of HTLV promoter; codA::upp, bacterial cytosine deaminase and uracil phosphoribosyltransferase conferring sensitivity to the drugs 5FU and 5-fluorocytosine.

Cells expressing the telomerase RNA gene will therefore activate the hTR promoter and cause expression of the HSV thymidine kinase gene, (Tk). Expression of the HSV Tk gene sensitises cells to the drug Ganciclovir. Thus, cells activating the hTR promoter in the suicide gene vector will be killed on exposure to Ganciclovir.

Materials and Methods

Site-directed mutagenesis and constructs:

The PCR mutagenesis were performed separately by using Site-directed mutagenesis kit [Stratagene, QuikChange™]. The primers used for the construction of site-replaced mutants

of the Sp1, CCAAT, RCE and TATA binding sites are shown in Figure 12.

The PCR reactions were performed using the following standard condition; All PCRs were performed using 2 μ l of
5 pC2923(-107/+69) plasmid DNA(50 ng/ μ l) as template, 1.5 mM MgCl₂, 0.2 mM dNTP's, 25 nM of site-replaced primer and 1.0 μ l(2.5 U/ μ l) Pfu DNA polymerase in 50 μ l reaction volume. A total of 16 cycles were used after originally denaturing at 95°C for 30 seconds. In each cycle, the protocol was as
10 follows:

30 seconds denaturing at 95 °C

30 seconds annealing at 55 °C

6 minutes elongation at 68 °C (2 minutes/Kb plasmid length)

15 Then, mutants of luciferase reporter construct were constructed by subcloning the mutation fragments into Xho I/Hind III linearized pGL3 Basic vector.

Annealing and gel purification of oligonucleotides

20 To anneal the two complementary oligonucleotides, the same molar ratio of oligonucleotides were mixed in TE buffer, the oligo was treated at 95 °C for 5 minutes in either the Perkin-Elmer Cetus DNA Thermal Cyclor or the Perkin-Elmer Cetus Cyclor 9600, then the Cyclor was
25 switched off to cool down slowly(about 4.5 hours). Then, the annealing oligo were loaded in the 12% PAGE gel and band excised. Finally, the pellet was resuspended in 25 μ l TE and quantitated.

Preparation of nuclear cell protein extracts

30 Nuclear protein preparation: Approximately 10¹⁰ cells

were washed in PBS and then twice with TMS (5 mM Tris-HCl pH 7.5, 2.5 mM MgCl₂, 125 mM sucrose). Cells were lysed in 200 ml of TMS plus 0.25% Triton X-100, and nuclei were harvested by centrifugation (1,600 x g, 20 minutes). Nuclei were washed three times in 200 ml of TMS and resuspended in approximately 5 ml of TMS (5 to 10 mg of DNA per ml), and 0.1 volumes of 4 M NaCl was added dropwise with stirring. The solution was centrifuged at 10,000 x g for 20 minutes and the supernatant was spun at 100,00 x g for 60 minutes. Solid ammonium sulfate was added to 0.35 g/ml and left on ice for 30 minutes. The precipitate was pelleted at 10,000 x g for 30 minutes and redissolved in 5 ml of E50 buffer (50 mM ammonium sulfate, 20 mM HEPES, pH 7.9, 5 mM MgCl₂, 0.1 mM EDTA, 0.1% [v/v] Brij 35, 20% [v/v] glycerol, 1 mM DTT) and dialyzed for 16 hours against 1 liter of storage buffer (50 mM NaCl, 20 mM HEPES, pH 7.9, 5 mM MgCl₂, 20% [v/v] glycerol, 1 mM DTT). The crude protein extract was cleared by centrifugation at 100.00 x g for 60 minutes, and aliquots were stored at -70°C.

Electrophoresis Mobility Gel Shift Assay (EMSA)

Electrophoresis Mobility Shift Assay (EMSA) was performed by EMSA kit [Promega].

Typical 5'-kinase labeling reactions included the DNA to be labeled, [gg-³²P]dATP [Amersham], T4 polynucleotide kinase, and buffer [Promega]. Synthesized oligomers were 5'-end-labeled with [gg³²P] dATP (5,000 mCi/mmol) and were used as probes in EMSA. 5.5 µg of HeLa nuclear proteins were incubated in 15 µl of reaction containing 4% glycerol, 1 mM MgCl₂, 0.5 mM dithiothreitol (DTT), 0.5 mM EDTA, 50 mM NaCl, 10 mM Tris-HCl, pH 7.5 and 2.0 µg poly(dI-dC), with or

without excess molar of unlabeled DNA competitors, on ice for 10 min, followed by addition of 5,000-10,000 cpm of the probe. For competition experiments, nonradioactive double-stranded oligodeoxynucleotides were added in 100-fold molar excess prior to addition of the probe. For supershift assays, rabbit anti-human polyclonal antibodies specific against Sp1, Sp3, Rb, Ap-2 (Santa Cruz Biotechnology) were added to the reaction mixture 25 minutes prior to the addition of the probe. All DNA-protein complexes were resolved on 5% native polyacrylamide gels run in 22.3 mM Tris, 22.3 mM boric acid, and 0.5 mM EDTA. Visualization was performed using both a computing PhosphorImager with ImageQuant software analysis (Molecular Dynamics) and autoradiography on Kodak XAR-5 films.

SuperFect™ Transfection (QIAGEN)

For co-transfection of wild type (pL2930,107/+69) and each individual mutation luciferase reporter construct, 3 ug plasmid DNA was mixed with 3 ug of Sp1, or Sp3 (pCMVSp1 and pCMVSp3), or Sp1 plus Sp3, or pCMVb expression vector, 0.25 ug pRB(pCMV-pRb) was used for co-transfection. 7.5 µl Superfect™ Transfection Reagent [QIAGEN] was added to the DNA solution and mixed by pipetting up and down 5 times to get complexes.

Discussion

In general, there are no significant sequence homologies between the promoter regions of the human and mouse telomerase RNA genes. Indeed, there is considerable debate as to whether telomere length is regulated in a similar fashion in humans and mouse (Blasco et al., 1997;

Kipling, 1997a; Kipling, 1997b; Zakian, 1997). However, mouse models represent a valuable resource with which to study the role of telomerase in cellular senescence and tumour progression and mouse models are likely to be required to investigate new therapies based on telomerase inhibition. In addition, the developmental regulation of telomerase will be more easily approached in mice (Bestilny et al., 1996; Blasco et al., 1995; Blasco et al., 1997; Blasco et al., 1996; Broccoli et al., 1996; Prowse & Greider, 1995). Thus, any differences between the two species may in fact aid understanding of the function for telomerase in maintaining genome stability and will be important in developing good murine models for human disease or developmental processes.

Despite the lack of sequence similarity between the human and mouse telomerase RNA gene promoter regions, they both have consensus sites for the binding of transcription factors implicated in haematopoiesis and leukaemogenesis such as GATA-1, PU.1, PEA2/PEBP2, C/EBP, and c-Ets-2 (Tenen et al., 1997). This data will therefore be used in the detection of telomerase activity in normal and malignant haematopoietic cells (Bodnar et al., 1996; Cheng et al., 1997; Holt et al., 1997; Norrback & Roos, 1997; Pan et al., 1997). The human and mouse telomerase RNA genes do share an interesting similarity, in that they both lie in CpG islands, and thus their expression may be regulated by methylation. DNA methylation is thought to be important for gene regulation during normal development and cellular senescence, and abnormal methylation patterns may be a fundamental change in tumour progression (Baylin et al., 1991; Bird, 1996; Laird & Jaenisch, 1996; Vertino et al.,

1994; Wilson & Jones, 1983). Thus it has been suggested that aberrant CpG island methylation during the normal ageing process, could contribute to immortalisation by interfering with expression of "mortality" genes, of which
5 hTR and *terc* can be included (Vertino et al., 1994; Wilson & Jones, 1983).

Turning to the functional analysis of the cloned sequences, the minimal promoter for hTR resides within a region of 272bp upstream of the published transcriptional
10 start site, (Feng et al., 1995), (Figures 5a, 4a). There are a number of potential transcription factor binding sites in this region including consensus sequences for AP1, Sp1, PEA2/PEBP2, PEA3 and PU.1. Interestingly, the expression in the fos/jun family of proteins, which
15 determine AP1 activity, are suppressed during the onset of senescence and would be predicted to lead to a reduction in AP1 activity in senescent cells (Campisi, 1997; Irving et al., 1992; Riabowol et al., 1992; Seshadri & Campisi, 1990). AP1 also responds to protein kinase C, and it has
20 recently been demonstrated that hTR expression is induced by protein kinase C during T-cell activation (Bodnar et al., 1996). Extending the promoter region to 463bp upstream of the transcriptional start site, increases the luciferase activity to its maximum level, (Figures 5a, 4a).
25 This region contains several consensus binding sites for glucocorticoid/progesterone/androgen receptor binding, which may contribute to the maximal activity demonstrated by hProm505. A reduction in promoter activity is observed on extending the promoter fragments to include more 5'-
30 sequence, (Figure 5a), suggesting that sequences towards

the 5'-end of the clone may influence promoter activity in a negative fashion.

The minimal promoter for *terc* resides in a 94bp region upstream of the published transcriptional start site, (Blasco et al., 1995), (Figures 5b, 4b). A striking feature of this region is the presence of three AP-2 consensus sites, two of which are coupled to c-Ets-2 sites and all these elements are contained in the 73bp region required for promoter activity, (Figures 5b, 4b). Oncogenic Ras gene signalling has been shown to operate through c-Ets-2 binding sites, thus there is a testable relationship between oncogene activation during tumour progression and telomerase RNA gene transcriptional activity (Galang et al., 1994; Wasylyk et al., 1994). A reduction in promoter activity is observed on extending the promoter fragments to include more 5'-sequence, (Figure 5b, mProm628), suggesting that sequences towards the 5'-end of the clone may influence promoter activity in a negative fashion.

In summary, the present inventor has provided the art with DNA sequence elements and gene products which regulate the hTR promoter. Thus, the inventor has identified the first direct molecular controls on signal transduction pathways affecting telomerase expression. The major findings are firstly, that the retinoblastoma gene product stimulates hTR promoter activity; secondly, that the Sp1 transcription factor stimulates hTR promoter activity; thirdly, that the Sp3 transcription factor is a potent repressor of hTR promoter activity; fourthly that mutation of the hTR CCAAT sequence elements abrogates promoter activity; and fifthly, that the hTR Sp1-1 sequence element is a site for a repressor of hTR promoter activity.

These studies show that alterations in the amounts or activities of Sp1/Sp3/pRb/CCAAT factors in a cell or signal transduction specific manner may contribute to hTR gene expression. Further, these studies provide evidence for the signal transduction pathways which may be operating in normal cells to regulate hTR gene expression. They provide a focus for studies designed to uncover how cancer cells reactivate or upregulate hTR gene expression. Transcription factors complexes on the hTR promoter as provided herein offer direct targets for developing inhibitory or gene-mimetic molecules, and assays for screening for them. The understanding of the molecular basis for hTR gene regulation will be of value in the manipulation of telomerase expression, for example, in age related disease or optimal extension of cellular lifespan of commercially useful cells. These studies further provide insight into the regulation of other components of the telomerase enzyme, for example, the protein component, hTERT.

References

- Avilion, A.A., Piatyszek, M.A., Gupta, J., Shay, J.W.,
Bacchetti, S. & Greider, C.W. (1996). *Cancer Research*,
56, 645-650.
- 5 Baylin, S.B., Makos, M., Wu, J., Yen, R.W., De, B.A.,
Vertino, P. & Nelkin, B.D. (1991). *Cancer Cells*, 3,
383-390.
- Benda, J.A. (1994). Semin. In *Oncol.*, 21, 3-11.
- Bestilny, L.J., Brown, C.B., Miura, Y., Robertson, L.D. &
10 Riabowol, K.T. (1996). *Cancer Research*, 56, 3796-3802.
- Bird, A.P. (1996). *Cancer Surveys*, 28.
- Blasco, M.A., Funk, W., Villeponteau, B. & Greider, C.W.
(1995). *Science*, 269, 1267-1270.
- Blasco, M.A., Lee, H., Hande, M.P., Samper, E., Lansdorp,
15 P.M., DePinho, R.A. & Greider, C.W. (1997). *Cell*, 91,
25-34.
- Blasco, M.A., Rizen, M., Greider, C.W. & Hanahan, D.
(1996). *Nature Genetics*, 12, 200-204.
- Bodnar, A.G., Kim, N.W., Effros, R.B. & Chiu, C.P. (1996).
20 *Experimental Cell Research*, 228, 58-64.
- Breslow, R.A., Shay, J.W., Gazdar, A.F.T. and Srivastava
S., (1997). *J. Natl. Cancer Inst.*, 89, 618-623.
- Broccoli, D., Chong, L., Oelmann, S., Fernald, A.A.,
Marziliano, N., Van, S.B., Kipling, D., Le, B.M.M. &
25 De, L.T. (1997). *Human Molecular Genetics*, 6, 69-76.
- Broccoli, D., Godley, L.A., Donehower, L.A., Varmus, H.E.
& De, L.T. (1996). *Molecular & Cellular Biology*, 16,
3765-3772.
- Bryan, T.M., Marusic, L., Bacchetti, S., Namba, M. &
30 Reddel, R.R. (1997). *Human Molecular Genetics*, 6, 921-
926.

- Burger, A. M., et al (1997). *Br. J. Cancer*, **75**, 516-522.
- Cai, W., Hu, L. & Foulkes, J.G. (1996). *Current Opinion in Biotechnology*, **7**, 608-615.
- Campisi, J. (1997). *European Journal of Cancer Part A*, **33**,
5 703-709.
- Cheng, A.J., Liao, S.K., Chow, S.E., Chen, J.K. & Wang, T.C. (1997). *Biochemical & Biophysical Research Communications*, **237**, 438-444.
- Connors, T.A. (1995). *Gene Therapy*, **2**, 702-709.
- 10 Feng, J., Funk, W.D., Wang, S.S., Weinrich, S.L., Avilion, A.A., Chiu, C.P., Adams, R.R., Chang, E., Allsopp, R.C., Yu, J., Le, S., West, M.D., Harley, C.B., Andrews, W.H., Greider, C.W. & Villeponteau, B. (1995). *Science*, **269**, 1236-1241.
- 15 Galang, C.K., Der, C.J. & Hauser, C.A. (1994). *Oncogene*, **9**, 2913-2921.
- Gardiner-Garden, M. & Frommer, M. (1987). *Journal of Molecular Biology*, **196**, 261-282.
- Harley, C.B. and Villeponteau, B. (1995), *Curr.Opin. Genet. Develop.*, **5**, 249-255.
- 20 Herwig, S. Et al (1997). *Eur.J. Biochem*, **246**, 581-601
- Hiyama, E., et al (1996). *J. Natl. Cancer Inst.*, **88**, 116-122.
- Hiyama, E., et al (1995a). *Nature Med.*, **1**, 249-255.
- 25 Hiyama, K., et al (1995b). *J. Nl. Cancer Inst.*, **87**, 895-902.
- Holt, S.E., Wright, W.E. & Shay, J.W. (1997). *European Journal of Cancer Part A*, **33**, 761-766.
- Holt, S.E., Shay, J.W. and Wright W.E. (1996a). *Nature Biotech.*, **14**, 836-839.
- 30 Holt, S.E., et al (1996b). *Mol. Cell. Biol.*, **16**, 2932-2939.

- Irving, J., Feng, J., Wistrom, C., Pikaart, M. & Villeponteau, B. (1992). *Experimental Cell Research*, **202**, 161-166.
- Kim, N.W. (1997). *European Journal of Cancer Part A*, **33**,
5 781-786.
- Kim, N.W., et al (1994) *Science* **266**, 2011-2015.
- Klingelhutz, A.J. et al (1996). *Nature*, **380**, 79-82.
- Kipling, D. (1997a). *Human Molecular Genetics*, **6**, 1999-2004.
- 10 Kipling, D. (1997b). *European Journal of Cancer Part A*, **33**, 792-800.
- Kuniyasu, H., Domen, T., Hamamoto, T., Yokozaki, H., Yasui, W., Tahara, H. & Tahara, E. (1997). *Japanese Journal of Cancer Research*, **88**, 103-107.
- 15 Laird, P.W. & Jaenisch, R. (1996). *Annual Review of Genetics*, **30**.
- Lania, L. et al (1997). *Int. J. Biochem. Cell Biol.* **29**, 1313-1323.
- Li, H., Zhao, L.L., Funder, J.W. & Liu, J.P. (1997).
20 *Journal of Biological Chemistry*, **272**, 16729-16732.
- Lundblad, V. and Wright, W.E. (1996). *Cell*, **87**, 369-375.
- Mandal, M. & Kumar, R. (1997). *Journal of Biological Chemistry*, **272**, 14183-14187.
- Miller, N. & Whelan, J. (1997). *Human Gene Therapy*, **8**, 803-
25 815.
- Morin, G.B. (1996). *Seminars in Cell and Developmental Biology*, **7**, 5-13.
- Morin, G.B. (1997). *European Journal of Cancer Part A*, **33**, 750-760.

- Nakamura, T.M., Morin, G.B., Chapman, K.B., Weinrich, S.L.,
Andrews, W.H., Lingner, J., Harley, C.B. & Cech, T.R.
(1997). *Science*, **277**, 955-959.
- 5 Norrback, K.F. & Roos, G. (1997). *European Journal of
Cancer Part A*, **33**, 774-780.
- Pan, C., Xue, B.H., Ellis, T.M., Peace, D.J. & Diaz, M.O.
(1997). *Experimental Cell Research*, **231**, 346-353.
- Parkinson, E.K. et al. (1997), *Euro. J. Cancer* **33**, 727-734.
- Peterson, M.G. & Baichwal, V.R. (1993). *Trends in*
10 *Biotechnology*, **11**, 11-18.
- Prowse, K.R. & Greider, C.W. (1995). *Proceedings of the
National Academy of Sciences of the United States of
America*, **92**, 4818-4822.
- Raymond, E. et al (1996) *Curr. Opin. Biothech.*, **7**, 583-591.
- 15 Riabowol, K., Schiff, J. & Gilman, M.Z. (1992). *Proceedings
of the National Academy of Sciences of the United
States of America*, **89**, 157-161.
- Seller, W.R. et al (1996). *Biochim. Biophys. Acta*, **1288**,
M1-5.
- 20 Seshadri, T. & Campisi, J. (1990). *Science*, **247**, 205-209.
- Sharma, S., Raymond, E., Soda, H., Sun, D., Hilsenbeck,
S.G., Sharma, A., Izbicka, E., Windle, B. & Von Hoff,
D.D. (1997). *Annals of Oncology*, **8**, 1063-1074.
- Shay, J.W. & Bacchetti, S. (1997). *European Journal of*
25 *Cancer Part A*, **33**, 787-791.
- Shay J.W. & Wright W.E. (1996). *Current Opinion in
Oncology* **8** 66-71.
- Soder, A.I., Goings, J.J., Kaye, S.B. & Keith, W.N. (1997a).
Oncogene, in press. - 16 979-983 (1998)
- 30 Soder, A.I., Hoare, S.F., Muir, S., Goings, J.J., Parkinson,
E.K. & Keith, W.N. (1997b). *Oncogene*, **14**, 1013-1021.

- Soder, A.I., Hoare, S.F., Muire, S., Balmain, A.,
Parkinson, E.K. & Keith, W.N. (1997c). *Genomics*, **41**,
293-294.
- 5 Tenen, D.G., Hromas, R., Licht, J.D. & Zhang, D.E. (1997).
Blood, **90**, 489-519.
- Vertino, P.M., Issa, J.P., Pereira-Smith, O.M. & Baylin,
S.B. (1994). *Cell Growth & Differentiation*, **5**, 1395-
1402.
- Villeponteau, B. (1996). *Seminars in Cell and*
10 *Developmental Biology* **7**, 15-21.
- Wasylyk, C., Maira, S.M., Sobieszczuk, P. & Wasylyk, B.
(1994). *Oncogene*, **9**, 3665-3673.
- Wellinger, R.J. & Sen, D. (1997). *European Journal of*
Cancer Part A, **33**, 735-749.
- 15 Wilson, V.L. & Jones, P.A. (1983). *Science*, **220**, 1055-1057.
- Wright, W.E., Piatyszek, M.A., Rainey, W.E., Byrd, W. &
Shay, J.W. (1996). *Developmental Genetics*, **18**, 173-
179.
- Yashima, K. Piatyszek M.A. Saboorian H.M. Virmani A.K.
20 Brown D. Shay J.W. and Gazdar A. F. (1997). *J. Clin.*
Path., **50**, 110-117.
- Zakian, V.A. (1997). *Cell*, **91**, 1-3.
- Zhao, J. Q. et al (1998) *Oncogene* **16** 1345-1350.

Claims

1. A nucleic acid molecule comprising a promoter
sequence derived from the telomerase RNA (TR) gene
5 promoter.
2. A nucleic acid molecule according to claim 1 wherein
the promoter sequence is derived from the human TR (hTR)
gene promoter.
- 10 3. A nucleic acid molecule according to claim 1 or
claim 2 wherein the promoter sequence is derived from a
nucleotide sequence as shown in Fig. 4a.
- 15 4. A nucleic acid molecule according to any one of the
preceding claims wherein the promoter sequence is at
least 230bp in length starting at position -42bp as shown
in Fig. 4a and Fig 5a upstream of the transcription start
site.
- 20 5. A nucleic acid molecule according to any of the
preceding claims wherein the promoter sequence comprises
the construct hProm505 as shown in Fig.4a and Fig.5a.
- 25 6. A nucleic acid molecule according to any one of the
preceding claims wherein the promoter sequence comprises
the construct hProm867 as shown in Fig. 4a and Fig. 5a.
- 30 7. A nucleic acid molecule according to any one of the
preceding claims wherein the promoter sequence comprises
the nucleotide sequence as shown in Fig. 4a.

8. A nucleic acid molecule according to claim 1 wherein the promoter sequence is derived from mouse TR (terc) gene promoter.

5 9. A nucleic acid molecule according to claim 8 wherein the promoter sequence is derived from the nucleotide sequence as shown in Fig. 4b.

10 10. A nucleic acid molecule according to claim 8 or claim 9 wherein the promoter sequence is at least 73bp in length starting at -22bp as shown in Fig. 4b and Fig. 5a upstream of the transcription start site.

15 11. A nucleic acid molecule according to any one of claims 8 to 10 wherein the promoter sequence comprises the construct mProm208 as shown in Fig. 4b and Fig. 5b.

20 12. A nucleic acid molecule according to any one of claims 8 to 11 wherein the promoter sequence comprises the construct mProm628 as shown in Fig. 4b and Fig. 5b.

25 13. A nucleic acid molecule according to any one of claims 8 to 12 wherein the promoter sequence comprises a nucleotide sequence as shown in Fig. 4b.

30 14. A nucleic acid molecule having promoter activity and being capable of hybridising under stringent conditions to the complementary sequence of a nucleic acid molecule according to any one of the preceding claims.

15. A nucleic acid construct comprising a TR promoter

region or a fragment, mutant, allele derivative or variant thereof able to promote transcription, operably linked to a heterologous gene.

5 16. A nucleic acid construct according to claim 15 wherein the heterologous gene encodes a cytotoxin.

10 17. A vector comprising a nucleic acid molecule according to any one of claims 1 to 14 or a nucleic acid construct according to claim 15 or claim 16.

15 18. A host cell comprising a nucleic acid molecule according to any one of claims 1 to 14 or a vector according to claim 17.

20 19. A host cell comprising a nucleic acid construct according to claim 15 or claim 16.

25 20. A method comprising culturing a host cell according to claim 19 under conditions for transcription of said heterologous sequence from the promoter.

30 21. A method according to claim 20 wherein the heterologous sequence is a coding sequence and the host cell is cultured under conditions for expression of the encoded peptide or polypeptide product.

35 22. A method according to claim 20 or claim 21 comprising detection of transcription of the heterologous sequence or the encoded product.

23. A method of screening for ability of a substance to modulate activity of a TR promoter, the method comprising contacting an expression system containing a nucleic acid construct according to claim 15 with a test or candidate
5 substance and determining transcription of the heterologous sequence or expression of the encoded peptide or polypeptide product.

24. A method according to claim 23 wherein the
10 expression system comprises a host cell containing said nucleic acid construct.

25. A method which comprises, following identification of a substance a having the ability to modulate activity
15 of a TR promoter in accordance with a method according to claim 23 or claim 24, manufacture of the substance and/or use of the substance in manufacture or formulation of a composition.

20 26. A substance having the ability to modulate activity of a TR promoter as determined by telomerase RNA gene expression.

25 27. A substance according to claim 26 which specifically blocks transcriptional activation of the TR gene promoter through interaction of the 5' regulatory sequences.

28. A substance according to claim 26 or claim 27 being
30 selected from the group consisting of antisense oligonucleotides, transcription factors, peptide nucleic acids, and factors that disrupt signal transduction

pathways.

29. A substance according to any one of claims 26 to 28
which is Sp3 transcription factor or the CCAAT protein
5 complex.

30. A substance according to any one of claims 26 to 29
for use in medical treatment.

10 31. Use of a substance according to any one of claims 26
to 29 in the preparation of a medicament for the
treatment of cancer.

32. A substance according to claim 26 which specifically
15 activates the TR gene promoter through interaction with
the 5' regulatory sequences.

33. A substance according to claim 32 which is Sp1
transcription factor or pRb transcription factor.

20 34. Use of a substance according to claims 32 or 33 for
the preparation of a medicament for the treatment of
cancer.

25 35. Use of a substance according to any one of claims 26
to 29 in the preparation of a medicament for repressing
the TR gene promoter in cells.

30 36. Use of a substance according to claims 32 or 33 for
the preparation of a medicament for activating the TR
gene promoter in cells.

37. A vector according to claim 17 for use in a method of gene therapy.

5 38. Use of a vector according to claim 17 in the preparation of a medicament for the treatment of cancer.

39. A system for use in the control of neoplasia in a human or animal subject comprising a vector or other delivery system capable of selectively infecting tumour
10 cells in said subject, said vector carrying a DNA or RNA sequence encoding an enzyme operably linked to a nucleic acid molecule according to any one of claims 1 to 14, in association with a prodrug capable of being converted to an active compound by action of said enzyme.

15

40. A system according to claim 39 wherein the vector is the gene therapy vector pGT62-codAupp.

20 41. A system according to claim 39 or 40 wherein the enzyme is viral thymidine kinase.

42. A system according to claim 39 or 41 wherein the prodrug is ganciclovir.

25 43. Use of a system according to any one of claims 39 to 42 in the preparation of a medicament for the treatment of neoplasia in a human or animal subject.

30 44. A method of treating neoplasia in a human or animal subject requiring such treatment comprising administering to the subject an effective amount of a prodrug and a

modified virus capable of selectively infecting tumour cells in said subject, said virus carrying a DNA or RNA sequence encoding an enzyme capable of converting said produg to an active compound, wherein said DNA or RNA
5 sequence is operably linked to a nucleic acid molecule according to any one of claim 1 to 14.

45. A method according to claim 44 wherein the enzyme is viral thymidine kinase and the prodrug is ganciclovir.

1/29

```

1  agtactcag gaggctgaga cagagaatc gcttgaaccc gggaggcaga ggtgacgtg
61  agccgagatc acgccactag actccatcca gcctggcgga aagagcaaga ctccgtctca
121  aaaaaaataa tcgttacaat ttatggtgga ttactcccc ttattacct catcaagaca
181  cagcactact ttaaaagcaa gtcaatgatt gaaacgcctt tctttcctaa taaaaggag
241  attcagtcct taagattaat aatgtagtag ttacacttga ttaaagccat cctctgctca
301  aggagaagct ggagaaggca ttctaaggaa aaaggggcag ggttgaact cggacgcac
361  cactgagcc gagacaagat tctgctgtag tcagtgtgc ctgggaatct atttcacaa
421  agttctccaa aaatgtgat gatcaaaact aggaattagt gtctgtgtc ttaggcccta
481  aaatcttct gtgaattcca tttttaaggt agtcgaggtg aaccgcgtct ggtctgcaga
541  ggatagaaaa aaggccctct gatacctcaa gttagtttca cctttaaga aggtcggaa
601  taaagacgca aagcctttcc cggacgtgcg gaagggaac gtccctctc atggccggaa
661  atggaacttt aatttcccgt tcccccaac cagcccgcc gagagagtga ctctcacgag
721  agccgcgaga gtcagcttgg ccaatccgtg cggtcggcg gagggtggt ggccattttt
781  ctgccccgc agcgacccgg gttcgggagg gtgggcctgg tgctccccg gcgtgtttt
841  tgtctaacc taactgagaa gggcgtagg ccgtgtctt tgcctccac gttcattcta
901  tctcgtgac tttcagcggg cggaaaagcc tggcctgcc ccgggggacc tgcggcgggt
961  gagcaaaaa aaaatgtcag ctgctggccc gttcgcccc cccgggggag ttctccggag
1021  cgcctgcca gccccgaac ccgcctgga ggcgcggtc ggcccgggc gggcgagagg
1081  gcacctactg ccaccgcgaa gatttggtc tgcagccgc cagtcctctg gggcgagagg
1141  cgagggttcag gcctttcagg ccgcaggaa aggaacggag cagtcctctg cgcgcggcgc
1201  gattccctga gctgtgggac gtgcacccag gactcggctc acacatgcag ttcgctttcc
1261  tgttggtgg ggaacgccg atcgtgcgca tccgtcacc ctcgcccga gtgggggctt
1321  gtgaacccc aaacctgact gactgggcca gtgtgctgca aattggcagg agacgtgaag
1381  gcacctccaa agtcgggcaa aatgaatgg cagtgaagcc gggctgctg gagccgttcc
1441  tgcgtgggtt ctcccgtctt ccgctttttg ttgctttta tgggtgtatt acaacttagt
1501  tctgctctg cagattttgt tgaggttttt gcttctccca aggtagatct cgaccagtcc
1561  cctcaacggg gtgtgggaga acagtcat ttttttgaga gatcatttaa catttaatga
1621  atatttaatt agaagatcta aatgaacatt ggaattgtg ttcccttaat ggtcatcggt
1681  ttatgccaga ggttagaagt tcttttttg aaaaattaga ccttggcgat gaccttgagc
1741  agtaggatat aacccccaca agctt

```

Fig. 1

2/29

```

1 aagcttggac ttgacaaaga aactgcagat catctggacc ccccccccc cccatttagg
61 tttaacaatg taccagctat ctgacttaag caaactgtgt tcctcataga taaggcggga
121 ctgctcatgg tcattgtgaa gttcagttgg gataaacaaa ttttaagggtg cataacaaaa
181 aacacaaaat gttggtgttt gtttaaaaaa aactaaagaa tttctggagg caggcagtta
241 cagaaaacat gctgatattc tgagttgcct gctagttggt gccattccac cagagtgaac
301 acatctctgt tgacctgat tttctgtagg tctgtctgtg tgtctgtcct ttctccagca
361 agggctgacc ctaatcgggg tcccaggacc caagccttga gaaaggcagt agtatgtcat
421 ctagttgaaa tgacacattc tctacagtgt ccaaagaca tctttgtgct agacagaaca
481 ttttattgga tggactatgg ctgaccactt ggcttggggg gggggggaag gggccgcca
541 gggcgggggt ccctcatttg cttgttatta acacttgctt gtttgtttac ttgttagtag
601 gaatctgctc taccacgtgg gttctacatg gttccacagg ggtcacctgg tccgtttttg
661 ttttctggga cagttttcac aaatgttgtc tagactccac gttggctttg aagcctacag
721 ctatgagcct ctgtgccagt ttatgcagta gtatctctcg ggttgcctt caccgttagt
781 agtgggtgctc ttagaaggca ccgtgatttt ttgctttcca tctctttccc ctgccatgcc
841 ttctgtggtt ctctgccagg caccaaactg ttcagaaact ctccagcccg gtagagaacg
901 gtagggggaa agaactgacg tgtggaaggg atgggcaggg agaagaggca ccgaactcgg
961 tcttaacaaa aaaaaaaaaa aaaaggagca ttagaaaaaa aaacaaattt gtgaccttga
1021 actacagacc tcctgcctca gcctcctaca agctgggatt ataggctcgg gtcagctacc
1081 cttgaaatct ttttctttct ggaactcagt acctggttgg ccatgcactc acaagagatc
1141 cgctgcctt ctgtctctca aattctggaa ttaaagattt gcgccacttt tccccacttc
1201 ccccccggc tgtgggagtg gactgggttg aagggtggaat tttttttttt tttttttttt
1261 tttagtga aaagggggga ttggaaatat ccctacttcc aactctagta tatttcagaa
1321 accaagcctc agagatgtgc gtgcgtgcgt gtgtgtgtgt gtatgtgtgt gtgtctcaca
1381 gcaagaaaca gattttatta tttatttttt atttatttat tttttgcaag tgactggcta
1441 ggaagagtgg ggaagcggga ggacaaatgg ggaagaggga gcatttccgc aagtgtcggg
1501 ctcgaccaat cagcgcgcgc catggggtat ttaaggtcga gggcggtag gcctcggcac
1561 ctaaccctga ttttcattag ctgtgggttc tggcttttg ttctccgccc gctgtttttc
1621 tcgctgactt ccagcgggccc aggaagtc agacctgcag cgggccaccg cgcgttcccg
1681 agcctcaaaa acaaacgtca gcgcaggagc tccaggttcg ccgggagctc cgcggcgccg
1741 ggccgcccag tcccgtaccc gcctacaggc cgcggcgccg ctggggtctt aggactccgc
1801 tgccgcccgc aagagctcgc ctctgtcagc cgcggggcgc cgggggctgg ggccaggccg
1861 ggcgagcgcc gcgaaggaca ggaatggaac tggccccgt gttcgtgtc ttacctgagc
1921 tgtgggaagt gcacccggaa ctcggttctc acaaccccca ttcccgtg ggaaatgccc
1981 cgctgcaggg cgggcccgtc gaacctgcga cttctgggga aaggggcttc ggtgtgagac
2041 ggtagccagc caaagggtat atatcgccct cagccccgt cccctccac ttttgtctaa
2101 tactcctgtt tctgttgtgc agattttgca ggcgtttcgc tggctctgcc tgaacgagct
2161 atgcagccat gtggctcctt ggggtggggg tgggatggg aggactacag gcgtagatct
2221 tcatactggg tttgtgtagt gctgggaatt gaacctagtt tcctaagttc tctatcaact

```

Fig. 2

3/29



```

2281 ggtattccca ttgtatggga gatttttttt ttcttttgta tatgggggcg ttgaacattt
2341 tgtaaacaaat tagaaaatct agtagttttt taatgaaaat gttcactttt ctttgtcttt
2401 gggatgcaaa acattacatt gaagctgaga agtttaaaga tgcgtgtctt cccctgccta
2461 ccttcgcggt cacacagaac ctgttatctt ttcagaaaag aaaatgagat aggcagggtg
2521 gatctggagt ttcaaggccc ttgcctggtc tgcagagtga gttaggccac accagaaaag
2581 tatgtgtcaa aaacaaagaa gaaaggcttt gtgggggggtg gggtagcaaa cgatcttaat
2641 cccctgtgct tgggaggccc gcaaggggga tatctgtgaa ggagacaaac aaagctacac
2701 tgtgtggtaa acaaaaaacg aggaggagga gcaagaagaa tatgagagcc cacggaagga
2761 agagtatcag tccccaggcc accagttcct caggggtaac tatgtttgtg agtgtctcgg
2821 tgccttgact tcctcagtac ttttctgggt tttagtcata aaaaacattg aagagatgaa
2881 gaagtgtatg tttagtaagt acataccaaa agtttgtgag ctatatgcat atagcaactc
2941 agtcacctga aacaggcccc ttgcagctaa catatttctt agtattacta ttataaagac
3001 taggggagtt tctaagccgg cactccttac aagggacgaa gccatgttca gctccagctt
3061 gccaagattc tgaaacccaa cgtcaagcct gacgagttcg agcctggcat ctctcagccg
3121 ctgctcgagc tggagatgac cacggatctc aaggcacagc tgtgggaact caacatcacc
3181 gaagccaagg aaaattgaag ttggtgggtg tcagaaggct gttataattt ttgtaccagt
3241 tcctcagctt aaatctttcc agaaaatcca agtctggcta gtttgtgaat tggagaaaaa
3301 gttcagcgga aagcacgtgt cttcattgct cagaagagga tctgtccaag ccaaccagga
3361 aaagctgtac gaaaaataag ccaaagcacc ctagaagctg caccctgaca gcagtgcattg
3421 tcttctcaag tgaaattgtg ggaaagagga tccatccgtg tgaaactgga tggcaatctg
3481 gagcaggttc atcttcctct ctggtacatc ccatgtctcc tcatctccat cctccccctt
3541 gcctctgtgt ctcatctcta aaactctcag cccatcttcc tttaccactg cccaatcaca
3601 ggctctagcc ttacctttca cctgccctca cctgcttata gacagcaatc tacatttctc
3661 cctttttgtc caattaaaag actcttttct ctcggatata aaatgagcac aactattatt
3721 accattctgt aattttataaa gtatagatag acctaacacc cagtctatca ttttgacagt
3781 taaataaagc attctgcaat cctatcctaa ctttaaaagg cttataattc tacacttggt
3841 atgtcctggc tcagcttgta tattagaaaa ccatctcaaa ttatatatat atatatatta
3901 cacacacaca tatgtatata tacatatata tgtatacaca cacacacata tatatatgta
3961 tatgtatgta tgtatgtata tatatatact tttaatgcta aatagcctgg gttggctaag
4021 actacttcaa tcctgccaga attc

```

Fig. 2 (cont)

4/29

*Fig. 3A**Fig. 3B*

hProm867
-798 agctactcaggaggctgagacacgagaatcgcttgaacccgggaggcaga
-748 ggttgacagtgagccgagatcacgccactagactccatccagcctgggcga
Zeste
-698 aagagcaagactccgtctcaaaaaaaaaaattcgttacaatttatggtgga
hProm697
-648 ttactccctctttttacctcatcaagacacagcactactttaagcaaa
GR
-598 gtcaatgattgaaacgcctttctttcctaataaaaggagattcagtcct
cMYB NF1 PEA3
-548 taagattaataatgtagtagttacacttgattaaagccatcctctgctca
AFP1/BRN2
-498 aggagaagctggagaaggcattctaaggaaaaaggggcagggttggaaact
PEA3/c-Eis-2 Sp1/NF-E2 cMYB
-448 cggacgcacccactgagccgagacaagattctgctgtagtcagtgtgc
Zeste GCN4/AP1
-398 ctgggaatctattttcacaaagtctctccaaaaaatgtgatgatcaaaact
myogenin GR GR
-348 aggaattagtggttctgtgtcttaggccctaaaaatcttctgtgaattcca
GR/PR/AR F2F/Pit-1a Pit-1a
-298 tttttaaggtagtcgaggtgaaccggtctctggtctgcagaggatagaaaa
hProm341 GATA-1
-248 aaggccctctgatacctcaagtttagtttcacctttaaagaaggtcggaag
E1A-F
-198 taaagacgcaaagccctttcccgagctgcggaagggaacgctccttcttc
NF1 PEA3 PU.1
-148 atggccggaaatggaactttaatttcccggttcccccaaccagcccgccc
Sp1
-98 gagagagtgcactctcacgagagccgcgagagctcagcttggccaatccgtg
API GCN4/AP1 CCAAT Box
-48 cggtcgagcgccgctccctttataagccgactcgcccggcagcgaccgg
hProm111 PEA2/PEBP2 GAGA TBP/TFIID
gttgcgagggtgggcctgggaggggtggtggccattttttgtctaaccc
hProm template
taactgagaagggcgta

Fig. 4A



6/29

mProm628
 -531 tgtgaccttgaactacagacctctgacctcagcctcctacaagctgggat
 PPAR/ELP GR
 -481 tataggctcgggtcagctacccttgaaatctttttctttctggaactcag
 H4TF-I
 -431 tacctggttgccatgcactcacaagagatccgcctgccttctgtctctc
 API/GCN4/Zeste SP-1
 -381 aaattctggaattaaagatttgcgccactttcccccactccacccccgg
 mProm458 C/EBPalpha, beta AP-2
 -331 ctgtgggagtggaactgggttgaaggtggaattttttttttttttttttt
 mProm418 p300
 -281 ttttagtgaaaaaaggggggattggaaatatccctactttcaactctagt
 CPI
 -231 atatttcagaaaccaagcctcagagatgtgctgctgctgctgtgtgtgtg
 mProm267
 -181 tgtatgtgtgtgtgtgtctcacagcaagaacagattttattatttttt
 GR F2F/Pit-1a
 -131 tattttatttttttttgcaagtgtgctaggaagagtgggggaagcggg
 mProm208 IRF-1,2 GCN4/AP1 AP-2 c-ETS-2
 -81 aggacaaatgggggaagaggagcatttccgcaagtgtctgggctcgaccaa
 mProm136 GR AP-2 c-ETS-2 AP-2 CCAAT Box/AP-1
 -31 tcagcgcgcccatgggggtatttaaggtcgaggcggttaggcctcggca

cctaaccctgattttcattagctgtgggttctgggtcttttgttctccgc
 template
 cgctgtttttctcgtgacttccagcgg mProm

Fig. 4B

7/29

*Fig. 5A**Fig. 5B*

8/29



Oligo's Used: human

Name	Sequence	Comments
hTR5	TACGCCCTTCTCAGTTAGGGTTAG	
hTR14	GGATCCTACGCCCTTCTCAGTTAGGGTTAG	hTR5 with BamHI site
hTR13F	ACTGAGCCGAGACAAGATTC	
hTR17F	GGATCCACTGAGCCGAGACAAGATTC	hTR13F with BamHI site
hTR10F	AGCTACTCAGGAGGCTGAGA	
hTR20F	GCGCTCGAGAGCTACTCAGGAGGCTGAGA	hTR10F with XhoI site plus gcg clamp
hTR11F	CATCAAGACACAGCACTACT	
hTR21F	GCGCTCGAGCATCAAGACACAGCACTACT	hTR11F with XhoI site plus gcg clamp
hTR6F	GTCTGGTCTGCAGAGGATAG	
hTR22F	GCGCTCGAGGTCTGGTCTGCAGAGGATAG	hTR6F with Xho site plus gcg clamp
hTR5	TACGCCCTTCTCAGTTAGGGTTAG	
hTR23R	CGCAAGCTTTACGCCCTTCTCAGTTAGGGTTAG	hTR5 with HindIII site plus cgc clamp
hTRe	CTGAGCTGTGGGACGTGCAC	
hTRf	AGACGGGAGAACCCACGCAG	
hTRg	CTCGGCTCACATGCAGTT	
hTRh	TCTGCAGAGCAGGAACTAAGT	
TRC3F	CTAACCCCTAACTGAGAAGGGCGTA	
TRC3R	GGCGAACGGGCCAGCAGCTGACATT	


Oligo's Used: Mouse

Name	Sequence	Comments
mTR16F	GTGTCTCACAGCAAGAAACA	
mtr25f	GCGCTCGAGGTGTCTCACAGCAAGAAACA	This is mtr16f with XhoI site plus gcg clamp
mTR17F	GTGACTGGCTAGGAAGAGTG	
mtr26f	GCGCTCGAGGTGACTGGCTAGGAAGAGTG	This is mtr17f with XhoI site plus gcg clamp
mTR18F	TGTGACCTTGAACCTACAGAC	
mtr27f	GCGCTCGAGTGTGACCTTGAACCTACAGAC	This is mtr18f with XhoI site plus gcg clamp
mTR19F	GGACTGGGTTGAAGGTGGAA	
mtr28f	GCGCTCGAGGGACTGGGTTGAAGGTGGAA	This is mtr19f with XhoI site plus gcg clamp
mTR20F	TGCGCCACTTTTCCCCACTT	
mtr29f	GCGCTCGAGTGCGCCACTTTTCCCCACTT	This is mtr20f with XhoI site plus gcg clamp
mTRr1	CCGCTGGAAGTCAGCGAGAA	
mtr30r	CGCAAGCTTCCGCTGGAAGTCAGCGAGAA	This is mTRr1 with HindIII site plus cgc clamp
mTR36F	GCGCTCGAGTCGACCAATCAGCGCGGCCAT	This is Xho I site PCR primer plus gcg clamp
mTRr1	CCGCTGGAAGTCAGCGAGAA	
mTRf1	TCGACCAATCAGCGCGGCCAT	

Fig. 6


**Fig. 7A****Fig. 7B**

10/29



tercCon

4044 bp

Fig. 8A

tercCon

4044 bp

Fig. 8B

11/29

TTTGTGACCTTGAACACTACAGACCTCCTGCCTCAGCCTCCTACAAGCTGGATTATAGGCTGGGTACGCTACCCCTTGAAA
TTCTTTTCTTTCTGGAACTCAGTACCTAGTGGCCATGCACACAGAGATCCGCCCTGCCCTCTGTCAAATTCCTGGA
ATTAAAGATTTCGCCACCTTTCCCCACTTCCACCCCCGGCTGTGGAGTGGACTGGGTGAAGGTGGAATTTTTTTTT
TTTTTTTTTTAGTGA AAAAGGGGGAATTGAAATATCCCTACTTTCAACTCTAGTATATTTCAAGAACCAAGCCTCAG
AAATGTGCGTGC GT
TTTATTTATTTTGTCAAGTACTGGCTAGGAAGAGTGGGGAACGGGAGGACAAATGGGGAAGAGGGAGCATTTCCGC
AAGTGTCTGGCTCGACCAATCAGCGCGGCCATGGGTATTTAAGTTCGAGGGCGGCTAGGCCCTCGGCACCTAACCCCTG
ATTTTCATTAGCTGTGGGTCTTCGCTTTCCTCCGCCCGCTGTTTTTCTCGCTGACTTCCAGGGGA

Fig. 9

12/29

Human Promoter

TATA

CCAAT

SP-1

API

PEA2/PEBP2

PU.1

PEA3


GATA1

C-MYB

GR/PR/AR

F2F

ZESTE



(540bp)

Fig. 10




Fig. 11

14/29

Oligonucleotide	Position	Sequence*	Purpose
h11 ^a	-2 to +36	CCGGTTCCGGAGGGTGGCTGGGAGGGTGGTGCC	RCE(+12,+16 and +30, +34) and Sp1.4 binding site
h111 ^a	"	CCGGTTCCGGAAATGGCTGGGAGGGTGGTGCC	RCE1 mutation from ggg to aaa (+11/+13)
h112 ^a	"	CCGGTTCCGGAGGGTGGCTGGGTAAGGTGGTGCC	Sp1.4 mutation from agg to taa (+24/+26)
h113 ^a	"	CCGGTTCCGGAAATGGCTGGGTAAGGTGGTGCC	Mutant of both RCE1 and Sp1.4 binding site(+11/+13, +24/+26)
h11c ^a	-2 to +23	CCGGTTCCGGAGGGTGGCTGGG	RCE1 binding site
h11d ^a	+14 to +36	CCCTGGAGGGTGGTGCC	Sp1.4 or RCE2 binding site
h111a ^{a,b}	"	CCGGTTCCGGAAATGGGCTGGG	RCE1 mutation from ggg to aaa(+11/+13)
h112b ^{a,b}	+15 to +36	GGCCTGGGTAAGGTGGTGCC	Sp1.4 mutation from agg to taa
h112c ^{a,b}	"	GGCCTGGGTAAGGTAATGGCC	Sp1.4 and RCE2 mutant from aggggtgg to taaggtaa(+24/+26, +30/+31)
h11e ^a	"	GGCCTGGGAGGGTAATGGCC	RCE2 mutant from gg to (+30/+31)
h10 ^a	-63 to -42	CTTGGCCAAATCCGTGCGTCCG	h10 footprinting region containing CCAAT binding site
h101 ^a	-63 to -42	CTTGGAGTCTCCGTGCGTCCG	CCAAT motif mutation from ccaa to agtc, (-58/-55)
h10m11 ^b	-74 to -45	GCAGAGTCAGCTTGGAGTCTCCGTGCGG	CCAAT motif mutation from ccaa to agtc, (-58/-55)
h10m2 ^{a,b}	-63 to -42	CTTGGCCAAATCCGTGAGTCCG	h10 mutation from gtcg to tgat, (-51/-47)
h9 ^a	-44 to -17	CGCGCGCGCTCCCTTTATAAGCGACT	h9 footprinting region containing SP1.1 binding site
h91 ^a	-44 to -21	CTTACGCGCTCCCTTTATAAGCC	h9 mutation from ggcg to ttac, (-43/-40)
h910 ^b	-53 to -29	CCGTGCGGTCTTACGCCGCTCCC	h9 mutation from ggcg to ttac, (-43/-40)
h911 ^a	-44 to -21	CGGGTAAACTCCCTTTATAAGCC	h9 mutation from ggcg to taaa, (-39/-36)
h92 ^a	-44 to -21	CGCGGCCATAGCCTTTATAAGCC	h9 mutation from gctc to atag, (-36/-33)
h921 ^a	-44 to -21	CGCGGCCGCTCATGCTATAAGCC	h9 mutation from cctt to atgc, (-32/-29)
h93 ^a	-44 to -21	CGCGGCCGCTCCCTTCGACAGCC	h9 mutation from tata to cgac, (-28/-25)
h930 ^b	-38 to -14	CCGCTCCCTTCGACAGCCGACTCGC	h9 mutation from tata to cgac, (-28/-25)
h4 ^a	-110 to -91	ACCAGCCCGCCGAGAGAGT	h4 footprinting region containing SP1.2 binding site
h41m ^a	-110 to -91	ACCAGCCCGAACGAGAGAGT	Sp1.2 mutation from cc to aa, (-101/-100)
h5 ^a	-471 to -452	GAAAAAGGGCAGGTTTGA	SP1.3 binding site
h5m ^a	-471 to -452	GAAAAAGGTTTCAGGGTTTGA	Sp1.3 mutation from gg to tt, (-463/-462)

* Nucleotides corresponding to promoter sequences are given in uppercase letters from 5' end to 3' end; specific nucleotides mentioned in Purpose column are underlined. Highlight indicate mutagenesis nucleotide.

^a Complementary lowerstrand sequence for EMSA not shown.

^b For PCR-directed in vitro mutagenesis, complementary lowerstrand sequence not shown.

Fig. 12

15/29




Fig. 13




Fig. 14




Fig. 15






Fig. 16




- Gel shift assays identify DNA/protein binding activity
- Do mutation of these binding sites influence the promoter activity?

19/29

*Fig. 17*

20/29

*Fig. 18*

21/29

```

-107                                     -55
2923(-107 +69)  AGCCCGCCCGAGAGAGTGACTCTCACGAGAGCCGCGAGAGTCAGCTTGGCCAA
29m23(mSp1.2)   -----AA-----
1011            -----AGTC-----
29m292(mSp1-1,2) -----AA-----
29m921(mSp1-1,2) -----AA-----

-54                                     -1
2923(-107 +69)  TCCGTGCGGTCGGCGGCCGCTCCCTTTATAAGCCGACTCGCCCGGCAGCGCACC
102             ---TGAT-----
910             -----TTAC-----
911             -----TAAA-----
92(mSp1.1)      -----ATAG-----
29m292(mSp1-1,2) -----ATAG-----
921(mSp1.1)     -----ATGC-----
29m921(mSp1-1,2) -----ATGC-----
930(mTATA)      -----CGAC-----
26n23(-51 +69)  -----

+1                                     +53
26n23(-51 +69)  GGGTTGCGGAGGGTGGGCCTGGGAGGGGTGGTGGCCATTTTTGTCTAACC
29111(mRCE)     -----AAA-----
29112(mSp1.4)   -----TAA-----
111(Sp1.4)*     ---//---AAA-----
112(RCE)*       ---//---TAA-----
113(mSP1+mRCE)* ---//---AAA-----TAA-----
115*            ---//---TAA---AA-----
114*            ---//---AAA-----TAA---AA-----

+54                                     +69
CTAACTGAGAAGGGCGTA

```

* Double mutagenesis by using pLh29m292 as template.

Fig. 19

22/29





Fig. 20

*Fig. 21*





Fig. 22

25/29







Fig. 24

27/29

*Fig. 25*

28/29




Fig. 26




Fig. 27

1/29

1 agctactcag gagctgaga cagagaatc gttgaacc gggaggcaga ggtgcagt
 61 agcggagatc acgccactag actccatcca gctgggcga aagagcaga ctccgtctca
 121 aaaaaaaa tcttacaat ttatgttggg tactccctt ctttttacct catcaagaca
 181 cagcactact ttaagcaaa gtcaatgatt gaacgcctt tctttcctaa taaaagggag
 241 attcagtctt taagattaat aatgtatgag ttacacttga ttaaagccat cctctgctca
 301 aggagaagct ggagaaggca ttctaaggaa aaaggggcag ggttggaaat cggacgcata
 361 ccactgagcc gagacaagt tctgctgtag tcagtgtgc ctgggaatct atttcacaa
 421 agttctccaa aaatgtgat gatcaaaat aggaattagt gttctgtctc ttaygcccta
 481 aaatcttct gtgaattcca tttttaeggt agtcgaggtg aaccgcgtct ggtctgcaga
 541 ggaagaaaa aagccctct gatactcaa gttagttcca ccttcaaga agtcggaa
 601 taagagcga agcctttcc cggacgtgc gaagggaac gtccttctc atggccgaa
 661 atggaacttt aatttccgt tcccccaac cagccgcct gagagagtga ctctcacgag
 721 agcgcgaga gtcagcttgg ccaatccgt cggcgcctgg cgggggtggt ggcattttt
 781 ctgcgccgac agccacccg gttgcggagg gtcggcctgg gcttccctt tgcctccgc
 841 gttctaaacc taactgaga gggcgtagg cccgtgcttt gcttccacc ggtcattcta
 901 tctcgtgac ttccagcgg ggaataagcc tggcctgcc cccggggacc tgcggcgggt
 961 gagcaacaa aaatgtcag ctgctggccc gttgcctct cccggggacc tgcggcgggt
 1021 cgcctgccc gccccggaac ccgcctgga ggcgcggtc ggcgcggggc ttctccggag
 1081 gcacctactg ccacgcgaa gatttgctc tgtcagccgc ggtctctctg ggggcgaggg
 1141 cagggttcag gctttccagg ccgaaggaa ggaacggag cagtcctccg cgcgcgggcg
 1201 gattccctga gctgtgggac gtgcaccag gactcggctc acacatgcag ttgccttcc
 1261 gttgtgtggg ggaacgcg atcgtgcga tccgtcacc ctgcgcggca gtgggggctt
 1321 gtgaacccc aaacctgact gactgggcca gttgtctgca aattggcagg agaagcgaag
 1381 gcaacctcaa agtcggcca aatgaatggg cagtggcgg ggttgcctg ggcctgtcc
 1441 tgcgtgggtt ctccgtctt ccgtttttt tgccttcta tggttgtatt acaacttagt
 1501 tctgtctcty cagattttgt tgggttttt gtttccca aggtagatct cgaaccagtc
 1561 ctcacacggg gttgtggaga acagtcatt tttttgaga gatcatttaa catttaata
 1621 atatttaatt agaagatcta aatgaacatt ggaattgtg ttcctttaat ggtcatcgg
 1681 ttatgccaga ggttagaagt ttctttttg aaaaattaga ccttggcgat gaccttgag
 1741 atagagatat aacccccaca agctt

Fig. 1

2/29

```

1  aagccttggaac ttgacaaaga aactgcagat catctggacc ccccccctccc cccatttagg
61  tttaacaaatg taaccagctat ctgacttaag caaactgtgt tcctcataga taaggcggga
121 ctgctcatgg tcattgtgaa gttcagttgg gataaacaaa ttttaagggtg cetaacaaaa
181 aacacaaaat gttggtgttt gtttaaaaaa aactaaagaa tttctggagg caggcagtte
241 cagaaaaaat gctgatatto tgagttgcoo gotagttggg gccattccac cagagtgaac
301 acatctctgt tgaccctgat tttctgtagg tctgtctgtg tgtctgtcct ttctccagca
361 agggctgacc ctaatcgggg tccaggagcc caagccttga gaagggcagt agtatgtcat
421 ctagttyaaa tgacacatcc totacagtg tccaaatgaca tctttgtgct agacagaaca
481 ttttatggga tggactatgg ctgaccactt ggcttggggg ggggggggag gggccggcaa
541 gggcgggggt cctcatttg cttgttatta acacttgcct gtttcttac tctgttagag
601 gaatctgctc taaccagtg gttctacatg gttccacagg ggtcacctgg tccgtttttg
661 tttctgagg cagttttcaa aaatgttgtc tagactccac gttggcttgg aagcctacag
721 ctatgagcct ctgtgccagt ttatgcagta gtatctctcg ggttgcctt caccgttagt
781 agtgggtgctc ttagaaggca ccgtgatttt ttgctttcca tctctttccc ctgccatgcc
841 ttctgtggtt ctgtgccagg caaaaaatg ttcagaacct ctccagcccg gttagagacg
901 gtagggggaa agaaatgacg tgtgggaagg atgggcaggg agaagaggca ccgaactcgg
961 tcttaaacaa aaaaaaaaaa aaaaaggagca ttgaaaaaaa aaacaaattt gtgaccttgc
1021 actacagacc tcttgctcca gctccttaca agctgggatt ataggctcgg gtcagctacc
1081 cttgaaatct ttttctttct ggaactcagt acctgggttg ccctgcactc acaagagatc
1141 cggctgcoct ctgtctctca aattctggaa ttaagattt gcgcacttt tccccacttc
1201 cccccccggc tgtgggagtg gaactgggtt aggttgggat tttttttttt tttttttttt
1261 tttagtgaaa aaaggcggga ttggaatat cctactttc aactctagta tatttcagaa
1321 accaagcctc agagatgtgc gtgctgtgct gtgtgtgtgt gtatgtgtgt gtgtctcaca
1381 gcaagaaaca gattttatta tttatttttt atttatttat ttttgcaag tgaactggta
1441 ggaagagtgg ggaagcggga ggaacaaatg ggaagaggga gcatttcggc aagtgttggg
1501 ctcgaccaat cagcgcgcgc catggggtat ttaaggtcga gggcggctag gctcggcac
1561 cttaacctga ttttcattag ctgtgggttc tggctctttg ttctcggccc gctgttttc
1621 tcgctgactt ccagcgggce aggaagtcce agacctgcag cgggcccagg cgcgttcccg
1681 agcctcaaaa acaaacgtca gcgcaggagc tccaggttcg ccgggagctc cgcggcggc
1741 ggccgcccag tcccgtaccc gctacaggcc cgggcgggc ctggggtctt aggactcggc
1801 tgcgcgcggc aagagctcgc ctctgtcagc cgcggggcgc cgggggctgg ggcaggccg
1861 ggcgagcggc gogaaggaca ggaatggaac tggctcccgt gttcgggtgc ttacctgagc
1921 tgtgggaagt gcaaccggaa ctgggttctc acaaccccca tcccgcctgg ggaatgccc
1981 cgtgacagg cgggcgcgta gaacctgcga cttctgggga aaggggcttc ggtgtgagac
2041 ggtagccagc caaagggtat atatgcctc cagccccgt cccctccac ttttgtctaa
2101 tactcctgtt tctgttgtgc agatttttga ggcgtttcgc tggctctgcc tgaacgagct
2161 atgcagccat gtggtccttg ggggtggggg tgggatggg aggaactacg gcgtagatct
2221 tcatactggg tttgtgtagt gctgggaatt gaacctagtt tcttsagttc tctatcaact

```

Fig. 2

3/29



```

2281 ggtattccca ttgtatggga gatttttttt ttcttttgta tatggggggc ttgaacattt
2341 tgtaaacaaat tagaaaatct agtagtcttt taatgaaaat gttcactttt ctttgtcttt
2401 gggatgcaaa acattacatt gaagctgaga agtttaaga tgcgtgtctt cccctgccta
2461 ccttcggtt cacacagaac ctgttatctt ttcagaaaag aaaatgagat aggcagggtg
2521 gatctggagt ttcaaggccc ttgootggtc tgcagagtga gttaggccac accagaaaag
2581 tatgtgtcaa aaacaaagaa gaaaggcttt gtgggggggtg gggtagcaaa cgatcttaat
2641 cccctgtgct tgggaggccc gcaaggggga tatctgtgaa ggagacaaac aaagctacac
2701 tgtgtggtaa acaaaaaacg aggaggagga gcaagaagaa tatgagagcc caaggaaagga
2761 agagtatcag tccccaggcc accagttcct caggggtaac tatgtttgtg agtgtctogg
2821 tgccttgact tctcagtac tttcttgggt tttagtcata aaaaacattg aagagatgaa
2881 gaagtgtatg tttagtaagt acataccaaa agtttgtgag ctatatgcat atagcaactc
2941 agtcacctga aacaggcccc ttgcagctaa catatttctt agtattacta ttataaagac
3001 taggggagtt tetaagcgg cactccttac aagggaacgaa gccatgttca gctccagctt
3061 gccagattc tgaaccccaa cgtcaagcct gacgagttcg agcctggcat ctctcagccg
3121 ctgctcgagc tggagatgac caggatctc aaggcacagc tgtgggaact caacatcacc
3181 gaagccaagg aaaattgaag ttggtggtgg tcagaaggct gttataattt ttgtaccagt
3241 tctcagctt aaatctttcc agaaaatcca agtctggcta gtttgtgaat tggagaaaaa
3301 gttcagcggg aagcacgtgt cttcattgct cagaagagga tctgtccaag ccaaccagga
3361 aaagctgtac gaaaaataag ccaagcacc ctagaagctg caccctgaca gcagtgcctg
3421 tcttctcaag tgaaattgtg ggaagagga tccatccgtg tgaaactgga tggcaactg
3481 gagcaggctt atcttctct ctggtacat ccatgtctcc tcatctccat cctccccctt
3541 gctctgtgt ctcatctct aaactctcag cccatcttcc tttaccactg cccaatcaca
3601 ggctctagcc ttacettcca cctgcccctc cctgcttata gacagcaatc tacatttctc
3661 cctttttgtc caattaaaag actcttttct ctggatata aatgagacc aactattatt
3721 accattctgt aatttataaa gtatagatag acctaacacc cagtctatca ttttgacagt
3781 taaataaagc attctgcaat cctatcctaa ctttaaaagg ctataatc tacacttggt
3841 atgtcctggt tcagcttgta tattagaaaa ccatctcaaa ttatatatat atatataatta
3901 cacacacaca tatgtatata tacatatata tgtatacaca cacacacata tatatatgta
3961 tatgtatgta tgtatgtata tatatatact ttaaatgcta aatagccctg gttggctaag
4021 actacttcaa tctgcagaa attc

```

Fig. 2 (cont)

4/29

*Fig. 3A**Fig. 3B*

5/29

†hProm867
 -798 agctactcaggaggctgagacacgagaatcgcttgaacccgggaggcaga
 -748 ggttgcagtgagccgagatcacgccactagactccatccagcctgggoga
 Zeste
 -698 aagagcaagactccgtctcaaaaaaaantcgttacaatttatggtgga
 -648 ttactccccctctttttaccctcatcaagacacagcactactttaagcaaa
 †hProm697
 GR
 -598 gtcaatgattgaaacgcctttcttttcttaataaaggagattcagtcct
 cMYB NFI PEA3
 -548 taagattaataatgttagttagttacacttgattaaagccatcctctgtctca
 APPI/BRN2
 -498 aggagaagctggagsaggcattctcaaggaaaaaggggcagggttggaaact
 PEA3/c-Eb-2 Sp1/NF-E2 cMYB
 -448 cgagccatccactgagccgagacaagattctgctgtagtcagtgctgc
 †hProm505
 Zeste GCN4/API
 -398 ctgggaatctattttcacaaagttctccaaaaaatgtgatcaaaact
 myogenin GR GR
 -348 aggaattagtggtctgtgtcttaggcctaaaaatcttctgtgaattcca
 GR/PAR F2F/P1-1a P1-1a
 -298 ctttaaggtagtcgaggtgaaccgctctgtgtctgcagaggatagaaaa
 †hProm341 GATA-1
 -248 aaggccctctgtatccctomagttagtttccctttaaagaaggtcggaaag
 E1A-F
 -198 ttaagacgcnaagcctttccgggaogtgoggaagggaaggtccttctc
 NFI PEA3 PU.1
 -148 atggccggaaatggaaactttaatttccgggtcccccnaaccagccccccc
 Sp1
 -98 gagagagtgaactatcaagagagccggagagagtcagcttggccaatccgtg
 API GCN4/API CCAAT Box
 -48 cggtccggccgctccctttataagccgactcgccgggcagcgaccgg
 †hProm111
 PEA2/PEBP2 OAOA TBP/TFIID
 gttgcggagggtgggcctgggaggggtggtggccatttttgtctaaacc
 template
 taactgagaaggcgta †hProm

Fig. 4A



6/29

mProm628
 -531 ttgtgaaccttgaaactacagacctcctgcctcagcctcctacagagctgggat
 PPAR/ELP GR
 -481 tataggctcgggtcagctacccttgaatatcttttctttctggaactcag
 H4TF-1
 -431 tacctggttggccatgcactcacaagagatccgcctgcctttctgtctctc
 API/GCN4/Zinc SP-1
 -381 aaattctggaattaaagatttgcgcacttttcccacttccacccccgg
 mProm438 C/EBPalpha, beta AP-2
 -331 ctgtgggagggactgggttgaagggtggaattttttttttttttttttt
 mProm418 p300
 -281 ttttagtgaaaaaaggggggttggaatatccctactttcaactctagt
 CPI
 -231 atatttcagaaaaccaagcctcagagatgtgggtgggtgggtgtgtgtgtg
 -181 tgtatgtgtgtgtgtgtctcagcagcaagaacagattttattatttttt
 mProm267 GR F2F/PK-1a
 -131 tatttatttatttttttgcagtgtactgggtaggaagagtggggaagcggg
 mProm208 (RF-1), 1 GCN4/AP1 AP-1 c-ETS-2
 -81 aggacaaatggggaagaggggagcaatttcaggcaagtgtgtgggtcgaccaa
 mProm136 GR AP-1 c-ETS-2 AP-2 CCAAT Box/AP-1
 -31 tcagcgcgcgccatgggggtatttaaggtcgagggggggtcaggcctcggca

ccataacctgattttcattagctgtgggttctggtcttttgtttctccgc
template
cgctgtttttctcgtgacttcacagcgg mProm

Fig. 4B

7/29

*Fig. 5A**Fig. 5B*

8/29



Oligo's Used: human

Name	Sequence	Comments
hTR5	TACGCCCTTCTCAGTTAGGXTTAA	
hTR14	GGATCCTACGCCCTTCTCAGTTAGGXTTAA	hTR5 with BamHI site
hTR13F	ACTGAGCCGAGACAAGATTG	
hTR17F	GGATCCACTGAGCCGAGACAAGATTG	hTR13F with BamHI site
hTR10F	AGCTACTCAGGAGGCTTACAA	
hTR20F	GCGCTCGAGAGCTACTCAGGAGGCTTACAA	hTR10F with XhoI site plus ggc clamp
hTR11F	CATCAAGACACAGCACTACT	
hTR21F	GCGCTCGAGCATCAAGACACAGCACTACT	hTR11F with XhoI site plus ggc clamp
hTR6F	GTCTGGTCTGCAGAGGATAA	
hTR22F	GCGCTCGAGGTCTGGTCTGCAGAGGATAA	hTR6F with XhoI site plus ggc clamp
hTR5	TACGCCCTTCTCAGTTAGGXTTAA	
hTR23R	CGCAAGCTTTACGCCCTTCTCAGTTAGGXTTAA	hTR5 with HindIII site plus cgc clamp
hTRc	CTGAGCTGTGGGACGTGCAC	
hTRf	AGACGGGAGAAACCCACGCAG	
hTRg	CTCGGCTCACACATGCAGTT	
hTRh	TCTGCAGAGCAGGAATAAGT	
TRC3F	CTAACCCCTAACTGAGAAAGGCGTA	
TRC3R	GGCGAAAGGGCCAGCAGCTGACATT	


Oligo's Used: Mouse

Name	Sequence	Comments
mTR16F	GTGTCTCACAGCAAGAAACA	
mtr25f	GCGCTCGAGGTGTCTCACAGCAAGAAACA	This is mtr16f with XhoI site plus ggc clamp
mTR17F	CTGACTGGCTAGGAAGAGTG	
mtr26f	GCGCTCGAGGTGACTGGCTAGGAAGAGTG	This is mtr17f with XhoI site plus ggc clamp
mTR18F	TGTGACCTTGAACCTACAGAC	
mtr27f	GCGCTCGAGTGTGAOCTTGAACCTACAGAC	This is mtr18f with XhoI site plus ggc clamp
mTR19F	GGACTGGGTGAAAGGTGGAA	
mtr28f	GCGCTCGAGGGACTGGGTGAAAGGTGGAA	This is mtr19f with XhoI site plus ggc clamp
mTR20F	TGGGOCACITTTCCCCACTT	
mtr29f	GCGCTCGAGTGGGCCACTTTTCCCCACTT	This is mtr20f with XhoI site plus ggc clamp
mTRr1	CCGCTGGAAGTCAGCGAGAA	
mtr30r	CGCAAGCTTCCGCTGGAAGTCAGCGAGAA	This is mTRr1 with HindIII site plus cgc clamp
mTR36F	GCGCTCGAGTGGACCAATCAGCGCGGCGCAT	This is XhoI site PCR primer plus ggc clamp
mTRr1	CCGCTGGAAGTCAGCGAGAA	
mTRr1	TGACCAATCAGCGCGGCGCAT	

Fig. 6


**Fig. 7A****Fig. 7B**

10/29



tarcCon

4044 bp

Fig. 8A

tarcCon

4044 bp

Fig. 8B

11/29

TTGTGACCTTGAACATACAGACCTCCTGCTCAGCCCTCCTACAAAGCTGGGATTATAGGCTCGGGTCAGCTACCCCTTGA
TTCCTTTTCCTTCTGGAACTCAGTACCTGGTTGGCCATGACATCACAAAGAGATCCGCCCTGCCCTTCTGTCAATTCCTGGA
ATTAAAGATTTGGCCACTTTTCCCACTTCCACCCCGGCTGTGGGAGTGGACTGGGTTGAAGGTGGAATTTTCTTTT
TTTTTTTATTAGTGAATAAGGGGGATTTGGAAATATCCCTACTTTCRACTCTAGTATATTTTCAGAAACCAAGCCCTCAG
AAATGTGCGTGCCTGCGTGTGTGTGTATGTGTGTGTGTCTCACAGCAAGAAACAGATTTTATTATTTTATT
TTTATTATTATTTTTCCTAAGTACTGGCTAGGAAGAGTGGGAGAGCGGGAGGACAAATGGGGAAGGGGAGCATTTCCGC
ARGTCTGGGCTCGAACCAATCAGCGCGGCCCATGGGTATTTAAGGTCTGAGGGCGGCTAGGCCCTCGGACCTTACCCCTG
ATTTTCATTAGCTGTGGGTTCTGGTCTTCGTTCTCCTDDECCCGCTGTTTTCCTGCTCACTTCCAGGGGA

Fig. 9

12/29

<i>Human Promoter</i>
TATA
CCAAT
SP-1
API
PEA2/PEBP2
PU.1
PEA3
GATA1
C-MYB
GR/PR/AR
E2F
ZESTE




Fig. 10

	-107	Sp1-2	
2923 (wt)		<u>AGCCCGCCGAGAGAGTGACTCTC</u> <u>ACGAGAGCCGCGAGAGTCAGCT</u> <u>TTGGCCAA</u>	CCAAT-box -55
	-54		
2923 (wt)		<u>TCCGTCCGGTCGGCGGCCGCTCCCTTTATAAGCCGACTCGCCCGGCGAGCCACC</u>	-1
	+1		
		Sp1-1	+52
		<u>GGGTTGGGAGGGGTGGGCCTGGGAGGGGTGGTGGCCATTTTGTCTAACC</u>	
		RCE	Sp1-4
	+53		+69
		CTAACTGAGAGGGCGTA	

Fig. 11

14/29

Oligonucleotide	Position	Sequence ^a	Purpose
n11 ^a	-2 to +36	CCGGCTTCGCGAGCGGTGCCTTCAACGCGATCAGCGGCC	RCE(+12, -16 and +30, +24) and Sp1.4 binding site
n111 ^a	"	CCGGCTTCGCGAGCGGTGCCTTCAACGCGATCAGCGGCC	RCE1 mutation from ggg to aaa (-11/+12)
n112 ^a	"	CCGGCTTCGCGAGCGGTGCCTTCAACGCGATCAGCGGCC	Sp1.4 mutation from agg to taa (+24/+25)
n113 ^a	"	CCGGCTTCGCGAGCGGTGCCTTCAACGCGATCAGCGGCC	Mutant of both RCE1 and Sp1.4 binding site(-11/+13, +24/+26)
n11c ^c	+2 to +23	CCGGCTTCGCGAGCGGTGCCTTCAACGCGATCAGCGGCC	RCE1 binding site
n11d ^d	+14 to +36	CCGGCTTCGCGAGCGGTGCCTTCAACGCGATCAGCGGCC	Sp1.4 or RCE2 binding site
n11ta ^{ab}	"	CCGGCTTCGCGAGCGGTGCCTTCAACGCGATCAGCGGCC	RCE1 mutation from ggg to aaa(+11/+12)
n11tb ^{ab}	+15 to +38	CCGGCTTCGCGAGCGGTGCCTTCAACGCGATCAGCGGCC	Sp1.4 mutation from agg to taa
n11tc ^{ab}	"	CCGGCTTCGCGAGCGGTGCCTTCAACGCGATCAGCGGCC	Sp1.4 and RCE2 mutant from aggggtagg to taaggtaa(+24/+26, +30/+31)
n11e ^a	"	CCGGCTTCGCGAGCGGTGCCTTCAACGCGATCAGCGGCC	RCE2 mutant from gg to (-20/+21)
n10 ⁱ	-83 to +42	CCTGGCGAATTCGCGGTGCCTTCAACGCGATCAGCGGCC	n10 bootstrapping region containing CCAAT binding site
n10t ^a	-63 to +42	CCTGGCGAATTCGCGGTGCCTTCAACGCGATCAGCGGCC	CCAAT motif mutation from ccaa to agac, (-58/-55)
n10m1 ^b	-74 to +45	CCAAAGGCGAATTCGCGGTGCCTTCAACGCGATCAGCGGCC	CCAAT motif mutation from ccaa to agac, (-59/-55)
n10m2 ^{ab}	-63 to +42	CCTGGCGAATTCGCGGTGCCTTCAACGCGATCAGCGGCC	n10 mutation from gggc to tgat, (-51/-47)
n9 ^j	-44 to -17	CCAGCGCGCGCTTCCTTTTAAGAAGCGAGT	n9 bootstrapping region containing Sp1.1 binding site
n9t ¹	-44 to -21	CCTTCCGCGCGCTTCCTTTTAAGAAGCGAGT	n9 mutation from gagc to tacc, (-42/-40)
n9t0 ^b	-53 to +29	CCAGCGCGCGCTTCCTTTTAAGAAGCGAGT	n9 mutation from gagc to tacc, (-42/-40)
n9t1 ^a	-44 to -21	CCAGCGCGCGCTTCCTTTTAAGAAGCGAGT	n9 mutation from gagc to taaa, (-39/-36)
n9t2 ¹	-44 to -21	CCAGCGCGCGCTTCCTTTTAAGAAGCGAGT	n9 mutation from gatc to aaag, (-36/-33)
n9t3 ¹	-44 to -21	CCAGCGCGCGCTTCCTTTTAAGAAGCGAGT	n9 mutation from catc to atgc, (-32/-29)
n9t4 ¹	-44 to -21	CCAGCGCGCGCTTCCTTTTAAGAAGCGAGT	n9 mutations from tata to cgaac, (-28/-25)
n9t0 ¹	-39 to +14	CCAGCGCGCGCTTCCTTTTAAGAAGCGAGT	n9 mutation from tata to cgaac, (-28/-25)
n4 ^k	-110 to -91	ACCAGCGCGCGCTTCCTTTTAAGAAGCGAGT	n4 bootstrapping region containing Sp1.2 binding site
n4tm ¹	-110 to -91	ACCAGCGCGCGCTTCCTTTTAAGAAGCGAGT	Sp1.2 mutation from cc to aa, (-104/-100)
n5 ^k	-471 to -462	GAAAAGCGCGCTTCCTTTTAAGAAGCGAGT	Sp1.3 binding site
n5m ^k	-471 to -462	GAAAAGCGCTTCCTTTTAAGAAGCGAGT	Sp1.3 mutation from gc to tc, (-463/-462)

* Nucleotides corresponding to promoter sequences are given in uppercase letters from 5' end to 3' end; specific nucleotides mentioned in Purpose column are underlined. Highlight indicate mutagenesis nucleotide.

Complementary powerstrand sequence for BMSA not shown.

For PCR-directed *in vitro* mutagenesis, complementary lower

² For PCR-directed *in vitro* mutagenesis, complementary low-strand sequence not shown.

Fig. 12





Fig. 14

17/29



h41 (-110 -91)

ACCAGCCCGCCGAGAGAGT

h41m(-101 -100)


:::::::::::AA::::::::::

Fig. 15




- Gel shift assays identify DNA/protein binding activity
- Do mutation of these binding sites influence the promoter activity?

19/29

*Fig. 17*

20/29



21/29

```

-107                                     -55
2923(-107 +69)  AGCCCGCCCGAGAGAGTGA CTCTCAGGAGAGCCCGCAGAGTCAGCTTGGCCCA
29m23(mSp1.2)  -----AA-----
1011                                     AGTC
29m292(mSp1-1,2) -----AA-----
29m921(mSp1-1,2) -----AA-----

-54                                     -1
2923(-107 +69)  TCCGTGCGGTCGGCGCGCCGCTCCCTTTATAAGCCGACTCGCCCGCCAGCCGACC
102             ---TGAT---
910             -----TTAC-----
911             -----TAAA-----
92(mSp1.1)      -----ATAG-----
29m292(mSp1-1,2) -----ATAG-----
921(mSp1.1)      -----ATGC-----
29m921(mSp1-1,2) -----ATGC-----
930(mTATA)      -----CGAC-----
26m23(-51 +69)  -----

+1                                     +53
26m23(-51 +69)  GGGTTGCGGAGGGGTGGGCGCTGGGAGGGGTGGTGGCCATTTTTTGTCTAACC
29111(mRCE)     -----AAA-----
29112(mSp1.4)   -----TAA-----
111(Sp1.4)*     ---//---AAA-----
112(RCE)*       ---//---TAA-----
113(mSp1+mRCE)* ---//---AAA-----TAA-----
115*            ---//---TAA---AA-----
114*            ---//---AAA-----TAA---AA-----

+54                                     +69
CTAACTGAGGAGGGCGTA

```

* Double mutagenesis by using pLh29m292 as template.

Fig. 19

22/29





Fig. 20

*Fig. 21*




Fig. 22

25/29




Fig. 23





Fig. 24

27/29

*Fig. 25*

28/29




Fig. 26




Fig. 27



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/11, 9/12, 15/85, C12Q 1/68	A3	(11) International Publication Number: WO 99/38964 (43) International Publication Date: 5 August 1999 (05.08.99)
(21) International Application Number: PCT/GB99/00308 (22) International Filing Date: 29 January 1999 (29.01.99) (30) Priority Data: 9801902.9 29 January 1998 (29.01.98) GB (71) Applicant (for all designated States except US): CANCER RESEARCH CAMPAIGN TECHNOLOGY LIMITED [GB/GB]; Cambridge House, 6-10 Cambridge Terrace, Regent's Park, London NW1 4JL (GB). (72) Inventor; and (75) Inventor/Applicant (for US only): KEITH, William, Nicol [GB/GB]; CRC Dept. of Medical Oncology, University of Glasgow, CRC Beatson Laboratories, Garscube Estate, Switchback Road, Glasgow G61 1BD (GB). (74) Agents: CRIPPS, Joanna, E. et al.; Mewburn Ellis, York House, 23 Kingsway, London WC2B 6HP (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 20 January 2000 (20.01.00)
(54) Title: PROMOTER REGIONS OF THE MOUSE AND HUMAN TELOMERASE RNA COMPONENT GENES (57) Abstract <p>The present invention relates to the identification of the genomic promoter region of the human and mouse telomerase RNA gene. Telomerase activity is necessary for the unrestricted proliferative capacity of many human cancers. It is proposed that mutation or dysregulation of the telomerase repression pathway may cause reactivation or upregulation of telomerase expression in cancer. The invention provides details of elements important for the regulation of telomerase RNA genes, including the Sp family of transcription factors. There is further provided methods for screening for elements having the ability for suppressing telomerase RNA gene promoter activity and use of such elements in the treatment of cancers. In addition, evidence is also provided for the development of new transcription based therapies for cancer and for genetic approaches to targeting therapeutic genes to cancer cells. Namely, (1) transcriptional repression and the disruption of signal transduction pathways regulating telomerase activation. (2) Tumour specific gene expression for genetic therapy via telomerase RNA gene promoters.</p> <div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;"> <p>A</p> </div> <div style="text-align: center;"> <p>B</p> </div> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/GB 99/00308

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/11 C12N9/12 C12N15/85 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 01835 A (ANDREWS WILLIAM H ;VILLEPONTEAU BRYANT (US); FUNK WALTER (US); FEN) 25 January 1996 (1996-01-25)	1-7, 14, 15, 17, 18, 23-28, 35, 37, 38
Y	page 24, line 13 -page 31, line 19 page 37, line 16 -page 43, line 13	15-21, 26-28, 30, 31, 35, 39-45
A	page 47, line 15 -page 55, line 19 page 65, line 9 -page 66, line 14 ----- -/--	26-38

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "Z" document member of the same patent family

Date of the actual completion of the international search

5 October 1999

Date of mailing of the international search report

19.10.1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Andres, S

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/GB 99/00308

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	PALMITER R D ET AL: "CELL LINEAGE ABLATION IN TRANSGENIC MICE BY CELL-SPECIFIC EXPRESSION OF A TOXIN GENE" CELL, vol. 50, 31 July 1987 (1987-07-31), pages 435-443, XP002055685 ISSN: 0092-8674 the whole document	15-21
Y	FENG J ET AL: "THE RNA COMPONENT OF HUMAN TELOMERASE" SCIENCE, vol. 269, no. 5228, 1 September 1995 (1995-09-01), pages 1236-1241, XP000645335 ISSN: 0036-8075 cited in the application	26-28, 30,31,35
A	the whole document	1-7
Y	RING, C. ET AL.: "Suicide gene expression induced in tumour cells transduced with recombinant adenoviral, retroviral and plasmid vectors containing the ERBB2 promoter" GENE THERAPY., vol. 3, 1996, pages 1094-1103, XP002117436 ISSN: 0969-7128 the whole document	39-45
X	WO 96 01614 A (ANDREWS WILLIAM H ;VILLEPONTEAU BRYANT (US); FUNK WALTER (US); GRE) 25 January 1996 (1996-01-25) cited in the application page 12, line 28 -page 13, line 25 figures 1,3 example 7	1-8,14, 17,18
X	HINKLEY, C. ET AL.: "The mouse telomerase RNA 5'-end lies just upstream of the telomerase template sequence" NUCLEIC ACIDS RESEARCH., vol. 26, 15 January 1998 (1998-01-15), pages 532-536, XP002106807 OXFORD UNIVERSITY PRESS, SURREY., GB ISSN: 0305-1048 the whole document	1-3,8,9, 14,17,18
X	PARKINSON E K ET AL: "The genetic basis of human keratinocyte immortalisation in squamous cell carcinoma development: the role of telomerase reactivation." EUROPEAN JOURNAL OF CANCER, (1997 APR) 33 (5) 727-34, XP002117437	26,31, 32,34, 36-38
A	page 731, paragraph 2 -page 732	26-38

-/--

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>LANDBERG, G. ET AL.: "Telomerase activity is associated with cell cycle deregulation in human breast cancer" CANCER RESEARCH., vol. 57, 1 February 1997 (1997-02-01), pages 549-554, XP002117438 ISSN: 0008-5472</p>	26-38
A	<p>HART L R : "TISSUE- SPECIFIC PROMOTERS IN TARGETING SYSTEMICALLY DELIVERED GENE-THERAPY" SEMINARS IN ONCOLOGY, (FEB 1996) VOL. 23, NO. 1, PP. 154-158., XP002117439 the whole document</p>	39-45
A	<p>WO 95 06486 A (OLDFIELD EDWARD J ;RAM ZVI (US); US HEALTH (US); BLAESE R MICHAEL) 9 March 1995 (1995-03-09) page 14, last paragraph -page 15, line 6 example 1 claims</p>	39-45
A	<p>HARLEY, C. & SHERWOOD, S.: "Telomerase, checkpoints and cancer" CANCER SURVEYS, vol. 29, 1997, pages 263-284, XP002117440</p>	
A	<p>GREIDER C W: "TELOMERE LENGTH REGULATION" ANNUAL REVIEW OF BIOCHEMISTRY, vol. 65, 1996, pages 337-365, XP002056801</p>	
P,X	<p>ZHAO, J.-Q. ET AL.: "Cloning and characterization of human and mouse telomerase RNA gene promoter sequences." ONCOGENE, vol. 16, 1998, pages 1345-1350, XP002106808 ISSN: 0950-9232 the whole document</p>	1-15,17, 18
P,X	<p>WO 98 11207 A (VILLEPONTEAU BRYANT ;HARLEY CALVIN (US); GERON CORP (US)) 19 March 1998 (1998-03-19) page 23, line 25 -page 32, line 13 page 33, line 27 -page 38</p>	1-7,15, 17-26, 30-32, 34-38

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB 99/00308

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 44 and 45 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

As a result of the prior review under R. 40.2(e) PCT,
no additional fees are to be refunded.

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☒ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/GB 99/00308

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Present claims 26, 27 and 32 relate to a substance defined by reference to a desirable characteristic or property, namely its ability to modulate the activity of a TR promoter.

The claims cover all substances having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such substances. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the substance by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the specific modulators as defined in claims 28, 29 and 33.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/GB 99/00308

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 2-7 (all totally) and claims 1,14,17-18, 37-38 (all partially)

A nucleic acid comprising a promoter region of the human telomerase RNA gene, constructs, vectors and host cells containing it.

2. Claims: 8-13 (all totally) and claims 1,14,17-18, 37-38 (all partially)

A nucleic acid comprising a promoter region of the mouse telomerase RNA gene, constructs, vectors and host cells containing it.

3. Claims: 16 (totally) and claims 15,17-22 (all partially)

A nucleic acid construct comprising a telomerase RNA gene promoter linked to a cytotoxin, and its use in the treatment of cancer.

4. Claims: 23-36 (all totally) and claims 15, 17-22 (all partially)

A method for screening for a substance being a modulator of the promoter of a telomerase RNA gene, the substances identified and their use in the treatment of cancer or the activation of telomerase.

5. Claims: 39-45

A system for controlling neoplasia.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/GB 99/00308

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9601835 A	25-01-1996	US 5583016 A	10-12-1996
		US 5776679 A	07-07-1998
		AU 696702 B	17-09-1998
		AU 2964795 A	09-02-1996
		AU 3095095 A	09-02-1996
		AU 9712998 A	18-03-1999
		BG 101103 A	31-10-1997
		BR 9508254 A	23-12-1997
		CA 2194393 A	25-01-1996
		CN 1158617 A	03-09-1997
		CN 1168696 A	24-12-1997
		CZ 9700034 A	15-10-1997
		EP 0778842 A	18-06-1997
		EP 0793719 A	10-09-1997
		FI 970026 A	03-03-1997
		HU 78054 A	28-07-1999
		JP 10505488 T	02-06-1998
		JP 10507067 T	14-07-1998
		NO 970041 A	06-03-1997
		NZ 289720 A	29-03-1999
		PL 318169 A	26-05-1997
		WO 9601614 A	25-01-1998
		US 5876979 A	02-03-1999
		US 5837857 A	17-11-1998
WO 9601614 A	25-01-1996	US 5583016 A	10-12-1996
		US 5876979 A	02-03-1999
		AU 696702 B	17-09-1998
		AU 2964795 A	09-02-1996
		AU 3095095 A	09-02-1996
		AU 9712998 A	18-03-1999
		BG 101103 A	31-10-1997
		BR 9508254 A	23-12-1997
		CA 2194393 A	25-01-1996
		CN 1158617 A	03-09-1997
		CN 1168696 A	24-12-1997
		CZ 9700034 A	15-10-1997
		EP 0778842 A	18-06-1997
		EP 0793719 A	10-09-1997
		FI 970026 A	03-03-1997
		HU 78054 A	28-07-1999
		JP 10505488 T	02-06-1998
		JP 10507067 T	14-07-1998
		NO 970041 A	06-03-1997
		NZ 289720 A	29-03-1999
		PL 318169 A	26-05-1997
		WO 9601835 A	25-01-1996
		US 5776679 A	07-07-1999
		US 5837857 A	17-11-1998
WO 9506486 A	09-03-1995	CA 2169557 A	09-03-1995
		EP 0716614 A	19-06-1996
		JP 9505557 T	03-06-1997
WO 9811207 A	19-03-1998	AU 4351997 A	02-04-1998

